

Exhibit 5

**UNITED STATES DISTRICT COURT
NORTHERN DISTRICT OF CALIFORNIA**

IN RE: ROUNDUP PRODUCTS
LIABILITY LITIGATION

MDL No. 2741

Case No. 16-md-02741-VC

This document relates to:
ALL ACTIONS

**EXPERT REPORT OF DR. CHARLES W. JAMESON, Ph.D.
IN SUPPORT OF GENERAL CAUSATION
ON BEHALF OF PLAINTIFFS**

Charles William Jameson, Ph.D.
Independent Consultant
May 12, 2017

Statement of Purpose

I have been asked to provide my expert opinions regarding the carcinogenic potential of glyphosate and glyphosate-based formulations. As a chemist and toxicologist, I evaluated the association of cancer, including non-Hodgkin's lymphoma ("NHL"), with exposure to glyphosate and/or glyphosate-based formulations. In performing my analysis, I relied on standard methods used in toxicology. I reviewed published, peer-reviewed scientific literature, publically available Government and Industry documents, and internal company documents and studies provided to me. All my opinions expressed in this report are based on a reasonable degree of scientific certainty. I reserve the right to supplement this report if additional information becomes available that are relevant to my opinions.

Qualifications

I am a private consultant in environmental toxicology specializing in carcinogenesis. I received my undergraduate degree in chemistry in 1970 from Mount Saint Mary's College, Emmitsburg, Maryland, and my Ph.D. in Organic Chemistry in 1975 from the University of Maryland, College Park. I started my career in 1965 where, as a rising high school senior, I spent the summer at a bioassay research laboratory first as a mouse room tech cleaning cages and later as an assistant in the chemistry lab mixing pesticides in rodent feed for the bioassay studies. Upon completion of my Ph.D. and a brief post-doc at the University of Maryland, I began working in 1976 as a contractor to the National Institutes of Health's (NIH) National Cancer Institute (NCI), serving as a senior chemist in support of NCI's Rodent Bioassay Program. In this capacity I was responsible for helping to monitor and evaluate the chemistry performed at the NCI's contract bioassay laboratories. In addition, I also provided support to the NCI staff for the identification of new substances to be studied in the NCI Bioassay Program. This

support included preparing and providing the background data from the scientific literature concerning exposure and the carcinogenic potential of the substance of interest. I was recruited by, and joined, the NCI in 1979 to serve as the chief chemist for their Rodent Bioassay Program and was responsible for directing and monitoring all chemistry activities, participating in the development of experimental protocols for the 2 year rodent bioassays conducted at the contract laboratories, and doing on-site inspections of all bioassay contract labs to insure they were following our protocols. In addition, I took over the responsibility as secretary for the NCI's Chemical Selection Working Group (CSWG) where I coordinated all activities for the identification of new substances to be studied in the Bioassay Program, including the oversight of the scientific literature searching, gathering and summarization process, documentation of the CSWG's review of the data and recommendations for study by the NCI, and the forwarding of the recommendation to the Director of the NCI Bioassay Program.

Following the formation of the National Toxicology Program (NTP) in 1978, the NCI Rodent Bioassay Program was transferred to the NIH's National Institute of Environmental Health Sciences (NIEHS) in 1980 where I transferred to and assumed the responsibility for all chemistry aspects of the NIEHS Division of Toxicology Research and Testing. I served as the program leader for chemistry in the National Toxicology Program (NTP) from 1978 until 1990. While chemistry program leader, I developed chemistry standards for bioassay studies that were widely accepted as an integral part of many toxicology-testing programs. I am listed as a contributor for the evaluation, interpretation and reporting of results for more than 100 chemicals studied in chronic two-year bioassay studies by the National Toxicology Program as published in the Technical Report Series (1980-1990). These bioassay studies were peer reviewed by the NTP Board of Scientific Counselors.

In 1990, I transferred to the NIEHS Director's Office and became involved with the NTP's Report on Carcinogens (RoC), working on it for more than 18 years, serving as its Director for 13 years before retiring from the NIEHS in February of 2008. The RoC is prepared in response to Section 301(b)(4) of the Public Health Service Act, which stipulates that the Secretary of the Department of Health and Human Services (DHHS) shall publish a report which contains a list of all substances which either are known or may reasonably be anticipated to be human carcinogens; and to which a significant

number of persons residing in the United States are exposed. This responsibility has been delegated by the Secretary to the Director, NTP. As Director of the RoC, I was responsible for the report's overall preparation, review and approval for the Director, NIEHS/NTP. In this capacity, I coordinated all review activities related to the RoC, which is one of the most visible and highly scrutinized activities of the NTP and the DHHS. I oversaw the identification and review of all new nominations for listing and delisting in upcoming editions of the RoC. I served as Chairman of the NIEHS RoC Review Committee, Chairman of the NTP Executive Committee's Interagency Working Group for the RoC, and Advisor to the NTP's Board of Scientific Counselors' Subcommittee for the RoC. I supervised the review of each nomination to the RoC, insuring all relevant information and data for each nomination was available for the review committees and managed the reviews by the three scientific review committees. Shortly after I became Director of the RoC in 1995, the Director, NTP, ordered that a review of the RoC be done to broaden input into its preparation, broaden the scope of scientific review associated with the Report, and provide review of the criteria used for inclusion of substances in the RoC. I coordinated this activity, which lead to revised criteria for the RoC being approved by the Secretary, DHHS in July of 1996. I served as Project Officer for the resource support contract for the preparation of the RoC, which included providing technical direction and coordination of the preparation of the documents prepared for each new nomination to the RoC as well as the preparation of 4 editions of the RoC for submission to the DHHS Secretary for approval.

I am the Senior Author for 69 NTP Report on Carcinogens Background Documents, which contained all available data concerning the exposure and potential carcinogenic activity of the substance being reviewed for possible listing in the RoC. I maintained a continuing liaison with other government agencies, private industries, other non-government research organizations and international organizations to keep abreast of work being done in chemical carcinogenesis, priorities for the listing of substances in the RoC, and resources available for the review of substances nominated for listing in the RoC. I served as the point of contact and focus for all RoC activities which included interacting with stakeholders from national and international government, industry, legal, consumer advocate, and other private concerns. I responded to requests for information from both the national and international press and private individuals on a

routine basis. Upon my retirement in 2008, I established CWJ Consulting LLC as a vehicle for providing expert consulting services in environmental toxicology specializing in carcinogenesis.

During my career, I participated as a Working Group Member for the United Nations' World Health Organization (WHO) International Agency for Research on Cancer (IARC). On several occasions, I served as either overall Chair of the Working Group or Chair of the Subgroup for Cancer in Experimental Animals evaluating cancer data and publishing monographs of the evaluation. I served as a consultant to the WHO, serving as a Task Group member to develop Environmental Health Criteria documents for partially halogenated chlorofluorocarbons (freons).

I am the author or co-author of over 80 peer reviewed scientific publications and nine book chapters. The vast majority of these publications relate to studies conducted in support of animal carcinogenesis bioassay programs. As mentioned above, I was the editor of four editions of the RoC, senior author for 69 NTP RoC Background Documents for substances reviewed for listing in the Report and listed as a contributor for the evaluation, interpretation and reporting of results for more than 100 chemicals studied in chronic two-year bioassay studies by the NTP as published in the Technical Report Series (1980-1990). I co-edited two books: "Chemistry for Toxicity Testing" and "Health and Safety for Toxicity Testing." A copy of my current curriculum vitae is attached as Exhibit A.

International Agency for Research on Cancer (IARC)

As an introduction, I would like to explain the International Agency for Research on Cancer's (IARC) review of glyphosate to assess its potential carcinogenicity, and the development of Monograph 112. The Working Group classified glyphosate as "probably carcinogenic to humans" (Group 2A) at their meeting in March of 2015. Following this meeting, there have been a number of publications (including, but not limited to, Williams et al.^{1, 2}; Chang and Delzell³, Solomon⁴) criticizing the IARC review process and conclusions.

The purpose of the *Monographs* is to render critical reviews and evaluations of carcinogenicity evidence of a wide range of human exposures.⁵ The *Monographs*

represent a hazard identification that involves examination of all relevant information to assess the strength of the available evidence that an agent can cause human cancer. Identifying carcinogens is a key step in cancer prevention, and this activity represents an important international activity towards improving public health. The IARC Preamble⁵ states that a “cancer ‘hazard’ is an agent that can cause cancer under some circumstances, while a cancer ‘risk’ is an estimate of the carcinogenic effects expected from exposure to a cancer hazard. The *Monographs* are an exercise in evaluating cancer hazards, despite the historical presence of the word ‘risks’ in the title. The distinction between hazard and risk is important, and the *Monographs* identify cancer hazards even when risks are very low at current exposure levels, because new uses or unforeseen exposures could engender risks that are significantly higher.” In other words, hazard assessment determines whether an agent can cause cancer.

For the review of glyphosate as it relates to Monograph 112, IARC performed a search for all relevant biological and epidemiological data from publically available sources and sent copies of the materials found to the Working Group participants approximately six months prior to the start of the meeting. In addition to the materials sent from IARC, Working Group participants perform their own independent search of the scientific literature. As the IARC Preamble notes, “with regard to epidemiological studies, cancer bioassays, and mechanistic and other relevant data, only reports that have been published or accepted for publication in the openly available scientific literature were reviewed.”⁵ IARC also considers relevant and publically available material from US Environmental Protection Agency (“EPA”). Studies determined to be irrelevant, inadequate, or published too late to be adequately evaluated were cited but were not summarized. This process of data collection is typical of all IARC *Monographs* and is the body of literature used by the Working Group participants during each Monograph analysis.

The IARC Working Group meeting takes places at its headquarters in Lyon, France and lasts for approximately seven to eight days, where the Working Group will then finalize the texts and formulate its final evaluations. Participants are assigned to one of four subgroups covering either exposure data, cancer in humans, cancer in experimental animals, or mechanistic and other relevant data. Working Group participants are also assigned individual chemicals or agents being evaluated and asked to prepare preliminary

working papers for their specific subgroup that are then distributed prior to the meeting. The subgroups prepare joint drafts and summaries in breakout sessions during the first few days. The entire Working Group meets in brief plenary sessions every day to get updates on the progress of each individual subgroup and to discuss any issues the subgroups may have identified. The final days of the meeting consists of plenary session meetings to discuss all relevant data, review the subgroup drafts and develop the final evaluations. The entire Monograph volume is considered the joint product of the Working Group, and there are no individually authored sections.⁵

For Monograph 112, I served as Chairman of the subgroup for Cancer in Experimental Animals to assess the carcinogenicity of several organophosphate pesticides that included glyphosate, the active ingredient in Roundup®. This meeting was held March 3-10, 2015 and the Working Group classified glyphosate as “probably carcinogenic to humans” (Group 2A). This classification was based on limited evidence in humans for the carcinogenicity of glyphosate where a positive association has been observed for NHL, sufficient evidence in experimental animals for the carcinogenicity of glyphosate and that mechanistic and other relevant data support the classification of glyphosate in Group 2A. To provide a better understanding of this, I will: discuss the process used by the Working Group to arrive at this classification, define terms, explain the types of evidence considered, explain the scientific criteria that guide the evaluations, and explain how conclusions were reached throughout the process.

The following summary of the Working Group’s evaluation of the available literature is offered here, but also found in the IARC’s Preamble⁵:

- Exposure Data: The Working Group concluded there is wide spread exposure to glyphosate based on its use as the active ingredient in Roundup® which is a broad-spectrum herbicide. Glyphosate is the most heavily used herbicide in the world⁶ and can be found in soil, air, surface water, groundwater, and food. According to several studies, glyphosate has also been detected in urine from persons around the world.⁷⁻¹⁰ The general population is mainly exposed to glyphosate through diet and from use as a household weed control.

- Cancer in Humans: The Working Group identified seven reports from the Agricultural Health Study (AHS) cohort and numerous reports from case-control studies

in the evaluation of the epidemiological studies reporting on cancer risks associated with exposure to glyphosate. This Working Group applied the Bradford Hill criteria in its analyses and determined that in several case–control studies there was an increased risks for NHL due to glyphosate exposure.¹¹⁻¹⁸ The Working Group further noted that the increased risk for NHL persisted in the studies that adjusted for exposure to other pesticides. The Working Group concluded a positive association has been observed for exposure to glyphosate and NHL and that there is “limited evidence” in humans for the carcinogenicity of glyphosate. IARC determines limited evidence of carcinogenicity for an agent when “a positive association has been observed between exposure to the agent and cancer for which a causal interpretation is considered by the Working Group to be credible, but chance, bias or confounding could not be ruled out with reasonable confidence.”⁵

•Cancer in Experimental Animals: The Working Group reviewed scientific literature and reports including two studies in which glyphosate was reported to be tested for carcinogenicity in male and female mice by dietary administration, five studies that tested glyphosate in male and female rats by dietary administration and in drinking-water in one study. Studies of a glyphosate-based formulation tested in drinking-water in one study in male and female rats and by skin application in one initiation–promotion study in male mice were also reviewed. They observed that in one feeding study in male CD-1 mice,¹⁹⁻²² glyphosate induced a positive trend in the incidence of kidney renal tubule carcinoma, a rare tumor in this strain of mice. A second feeding study²³ reported a positive trend for hemangiosarcoma (a blood vessel tumor) in male mice. Glyphosate also increased pancreatic islet-cell adenoma in male rats in two feeding studies.²⁴⁻²⁶ The Working Group concluded there is “sufficient evidence” in experimental animals for the carcinogenicity of glyphosate. IARC defines “sufficient evidence” in experimental animals is as “a causal relationship has been established between the agent and an increased incidence of malignant neoplasms or of an appropriate combination of benign and malignant neoplasms in (a) two or more species of animals or (b) two or more independent studies in one species carried out at different times or in different laboratories or under different protocols.”⁵

•Mechanistic and Other Relevant Data: The Working Group reported the mechanistic data literature contained strong evidence that glyphosate causes genotoxicity

and oxidative stress. The strong evidence of genotoxicity came from studies conducted in human cells in vitro,²⁷⁻³² in mammalian model systems in vivo^{27,32} and in vitro,^{33,34} and from studies in other non-mammalian organisms^{29,35,36,37}, all of which yielded largely positive results. The Working Group also found strong evidence for genotoxicity caused by glyphosate-based formulations. There were three studies of genotoxicity end-points in community residents exposed to glyphosate-based formulations, two of which reported positive associations.^{38,39} Strong evidence for oxidative stress was determined by studies conducted in human cells in vitro^{28,40,41} and in many rodent tissues in vivo.^{32,42,43} The Working Group found weak evidence that glyphosate or glyphosate-based formulations induce receptor-mediated effects,^{44,45} may affect cell proliferation or death,^{44,46} and may also affect the immune system in rodents⁴⁷ and fish.^{48,49} The Working Group considered the body of evidence described above as a whole and reached an overall evaluation of Group 2A: glyphosate is probably carcinogenic to humans. IARC uses this category when evidence of carcinogenicity in humans is limited and evidence of carcinogenicity in experimental animals is sufficient.⁵

IARC uses the hazard identification process for its review, and this was done for Monograph 112. Hazard identification reflects the toxicological “law” of specificity of effects⁵⁰. Hazard identification uses a strength of the evidence approach. As applied, the Working Groups for Monograph 112 rigorously assessed the toxicological, mechanistic, and epidemiological data to form a judgment regarding the likelihood that the agent produces cancer.

Information Reviewed

During the course of work on this case, I reviewed the following materials:

- scientific literature relating to the carcinogenicity of glyphosate and/or glyphosate-based formulations;
- government documents relevant to assessing the carcinogenic hazard and risks associated with glyphosate and/or glyphosate-based formulations; and,
- various studies and documents produced in the litigation.

For a list of additional materials I reviewed, please see Exhibit B.

Description of the Methodology Used to Assess Carcinogenic Potential Associated with Exposure to Glyphosate and/or Glyphosate-Based Formulations.

Toxicologists routinely assess the hazards to human health related to exposure to chemicals in the everyday environment using a process called hazard identification. A hazard is any agent that can cause harm or damage to humans, property, or the environment.⁵¹ In other words, a hazard is any agent that can cause a specific damage. In this case, the hazard being examined is glyphosate and/or glyphosate-based formulations, the specific damage is NHL, and the hazard assessment I am making is to determine whether or not glyphosate and/or glyphosate-based formulations can cause NHL. The terms hazard and risk are often used interchangeably; however, these are two distinct terms. Risk is defined as the probability that exposure to a hazard will lead to a negative consequence, or more simply, $\text{risk} = \text{hazard} \times \text{dose (exposure)}$.⁵²

Toxicology is the basis on which hazard identification is established. Hazard assessment has been used for over four decades by a wide variety of governmental and nongovernmental organizations to evaluate the potential adverse health effects from chemical exposures. Hazard identification is a standard tool used by toxicologists when they are trying to determine if exposure to a chemical(s) can cause an adverse health effect in humans and is the first step in risk analysis. Hazard identification is performed by identifying the chemical someone has been exposed to and then reviewing the available toxicity data to outline the spectrum of adverse effects that would be associated with exposure to that particular chemical.⁵³ The toxicity data could be from studies in humans, in whole animals, or in cells, or could be data collected on chemically-similar substances when data on the chemical of interest are limited.

I used the following criteria for my hazard based assessment of glyphosate and/or glyphosate-based formulations, that is based on the criteria I developed for the Report on Carcinogens⁵⁴ and is the same as defined and characterized by IARC⁵:

- Cancer in Humans – Numerous case-control studies and the Agricultural Health Study (AHS) cohort reporting on possible associations of cancer and exposure to glyphosate were evaluated for any evidence of a causal relationship between glyphosate and human cancer.

- “Sufficient” evidence is defined as when a causal relationship was established between exposure to glyphosate and cancer and that chance, bias and confounding could be ruled out.⁵
 - “Limited” evidence is defined as a positive association has been observed between exposure to glyphosate and cancer and a causal interpretation is credible but alternative explanations such as chance, bias or confounding could not be ruled out.⁵
 - “Inadequate” evidence is defined as available studies are of insufficient quality, consistency or statistical power to permit a conclusion regarding a causal association between glyphosate exposure and cancer.⁵
- Cancer in Experimental Animals – the experimental animal studies reporting on possible associations of cancer and exposure to glyphosate were evaluated for any evidence of a causal relationship between glyphosate and cancer.
- “Sufficient” evidence is defined as a causal relationship between exposure to glyphosate and an increased incidence of malignant and/or a combination of malignant and benign tumors, in multiple species or at multiple tissue sites or from multiple studies, or by multiple routes of exposure, or to an unusual degree with regard to incidence, site, or type of tumor, or age at onset.⁵
 - “Limited” evidence is defined as the data suggest a carcinogenic effect but are limited for making a definitive evaluation because, e.g. the evidence of carcinogenicity is restricted to a single experiment; there are unresolved questions regarding the adequacy of the design, conduct or interpretation of the studies; or the agent increases the incidence only of benign neoplasms or lesions of uncertain neoplastic potentials.⁵
 - “Inadequate” evidence is defined as studies that cannot be interpreted to show either the presence or absence of a positive carcinogenic effect because of major qualitative or quantitative limitations such as inadequate numbers of animals, lack of adequate pathology, poor survival, major impurities in the test agent, too low a dose to see an effect, etc. It should be noted that although animal testing is routinely used to identify cancer hazard, the sites

of cancer observed in animals do not always correlate directly with the sites of cancer that would be observed in humans⁵⁵. This can be due to the differences in metabolism in laboratory animals and humans, differences in pharmacokinetics, or differences in tissue reactivity (pharmacodynamics) between species. Animal studies, instead, are used to identify a threat of cancer that is applied to human health hazard assessment⁵⁵. All chemicals known to induce cancer in humans, that have been studied under adequate experimental conditions, also cause cancer in laboratory animals⁵⁵ and underscores the concept that chemicals found to cause cancer in laboratory animals must be considered capable of causing cancer in humans.⁵

- Mechanistic and other data – studies containing data relevant to the possible mechanism(s) of glyphosate carcinogenesis (genetic toxicity, epigenetic effects, etc.) were also evaluated. Mechanistic data may provide evidence of carcinogenicity and help in assessing the relevance and importance of findings of cancer in animals and humans.⁵

Hazard Assessment of the Human Data for Glyphosate and/or Glyphosate-Based Formulations

Before discussing the human data for glyphosate and/or glyphosate-based formulations, I will define the type of epidemiology studies that were reviewed:

- Case-Control Study - In a case-control study, investigators start by enrolling a group of people with disease. As a comparison group, the investigator then enrolls a group of people without disease (controls). Investigators then compare previous exposures between the two groups. The control group provides an estimate of the baseline or expected amount of exposure in that population. If the amount of exposure among the case group is substantially higher than the amount you would expect based on the control group, then illness is said to be associated with that exposure. The key in a case-control study is to identify an appropriate control group, comparable to the case group in most respects, to provide a reasonable estimate of the baseline or expected exposure.⁵⁶

- Cohort Study - According to Centers for Disease Control and Prevention (CDC),⁵⁷ in a cohort study the epidemiologist records whether each study participant is exposed or not, and then tracks the participants to see if they develop the disease of interest. After a

period, the investigator compares the disease rate in the exposed group with the disease rate in the unexposed group. The unexposed group serves as the comparison group or control, providing an estimate of the baseline or expected amount of disease occurrence in the community. If the disease rate is substantively different in the exposed group compared to the unexposed group, the exposure is said to be associated with illness.

- **Meta-Analysis** – A meta-analysis is an important component of systematic review procedure that combines and analyzes quantitative and qualitative data from several separate but similar experiments or studies to test the pooled data for statistical significance. Combining the results of multiple studies produces a weighted average of the included study results and leads to a conclusion with greater statistical power and point estimate than would be possible from any individual study.

Case Control Studies

- Cantor et al. (1992)¹⁴ evaluated the incidence of NHL among males located in Iowa and Minnesota. A total of 622 men and 1245 population-based controls were included in the study. The association with farming occupation and specific agricultural exposures were evaluated. When compared with non-farmers, the positive associations (odds ratios) for NHL were significant at 1.2 (95% CI, 1.0–1.5) for men who had ever farmed, and not significant at 1.1 (95% CI, 0.7–1.9) for 26 exposed cases for ever handling glyphosate and adjusted for confounders (vital status, age, state, cigarette smoking status, family history of lymphohaematopoietic cancer, high-risk occupations, and high-risk exposures).

- DeRoos et al. (2003)¹¹ pooled the data from three case–control studies^{12–14} to study pesticide exposures as risk factors for NHL in men. Of a total study population of 870 cases and 2569 controls, there were 650 cases and 1933 controls included for the analysis of 47 pesticides that also controlled for potential confounding by other pesticides. A positive association (odds ratios) for the association between exposure to glyphosate and NHL in the 36 cases exposed was reported to be significant at 2.1 (95% CI, 1.1–4.0) in the logistic regression analyses but not in the hierarchical regression analysis (which uses a more conservative adjustment estimate) at 1.6 (95% CI, 0.9–2.8).

- The effect of asthma as a modifier of the association between pesticide exposure and NHL was reported on by Lee et al. (2004)⁵⁸. The study contained 872 cases diagnosed

with NHL, 45 of which had been told they also had asthma and 2381 matched controls, 132 reporting to have asthma. Individuals in the study group with a history of asthma had a non-significantly lower risk of NHL than non-asthmatics and no effect was seen with pesticide exposure. A positive associations (odds ratio) for NHL associated with glyphosate use were reported but were not significant at 1.4 (95% CI, 0.98–2.1; 53 exposed cases) among non-asthmatics and 1.2 (95% CI, 0.4–3.3; 6 exposed cases) for asthmatics, when compared with non-asthmatic non-exposed farmers.

- The associations between exposure to pesticides and NHL was studied by McDuffie et al. (2001)⁴⁵ in a multicenter population-based study that included 517 cases and 1506 controls among men of six Canadian provinces. A non-significant positive association (odds ratios) of 1.26 (95% CI, 0.87–1.80; 51 exposed cases; adjusted for age and province) and 1.20 (95% CI, 0.83–1.74, adjusted for age, province, high-risk exposures) were observed for exposure to glyphosate. In an analysis by frequency of exposure to glyphosate, participants with more than 2 days of exposure per year had a statistically significant positive association (odds ratio) of 2.12 (95% CI, 1.20–3.73, 23 exposed cases) compared with those with some, but less than 2 days of exposure.

- Nordstrom et al (1998)⁵⁹ conducted a study in Sweden on hairy cell leukemia (considered to be a subtype of NHL). There were 121 cases in men and 484 controls matched for age and sex. A non-significant age-adjusted positive association (odds ratio) of 3.1 (95% CI, 0.8–12; 4 exposed cases) was reported for exposure to glyphosate.

- Hardell and Eriksson (1999)⁶⁰ reported on the results of the incidence of NHL in men associated with pesticide exposure in four northern counties in Sweden and included 404 cases and 741 controls. The authors reported a non-significant positive association (odds ratio) for ever-use of glyphosate of 2.3 (95% CI, 0.4–13; 4 exposed cases) in an analysis of glyphosate only, and 5.8 (95% CI, 0.6–54) in a multivariable analysis.

- Hardell et al. (2002)¹⁷ performed a pooled analysis of two case-control studies, one on NHL⁶⁰ and another on hairy cell leukemia.⁵⁹ These pooled analyses were based on 515 cases and 1141 controls. A significant positive association was found for exposure to glyphosate compared to controls (odds ratio, 3.04; 95% CI, 1.08–8.52; 8 exposed cases), but the positive association (odds ratio) decreased to a non-significant 1.85 (95% CI, 0.55–6.20) when study area, and vital status were considered.

•A large population based case–control study of exposure to pesticides as a risk factor for NHL in Sweden was conducted by Eriksson et al. (2008)¹⁸. There were 910 cases and 1016 controls included in the study. The association (odds ratio) for exposure to glyphosate to NHL was positive and significant at 2.02 (95% CI, 1.10–3.71) compared to controls, but positive and non-significant at 1.51 (95% CI, 0.77–2.94) when confounders that included exposure to other pesticides, age, sex, and year of diagnosis or enrolment were included in the analysis. When exposure to glyphosate for more than 10 days per year was considered, the positive association (odds ratio) was significant at 2.36 (95% CI, 1.04–5.37). Considering a latency period of greater than 10 years gave a positive association (odds ratio) that was also significant at 2.26 (95% CI, 1.16–4.40). The authors also reported an association with exposure to glyphosate and lymphoma subtypes. Positive associations were reported for most of the cancer forms, including B-cell lymphoma (odds ratio of 1.87; 95% CI, 0.998–3.51, non-significant) and the subcategory of small lymphocytic lymphoma/chronic lymphocytic leukemia (odds ratio of 3.35; 95% CI, 1.42–7.89, significant). These odds ratios were not adjusted for other pesticides.

•Orsi et al. (2009)⁶¹ reported the results of a case–control study conducted in France. The study included 491 cases (244 cases of NHL, 87 cases of Hodgkin lymphoma, 104 of lymphoproliferative syndrome, and 56 cases of multiple myeloma), and 456 age- and sex-matched controls. Positive, non-significant associations (odds ratios) for any exposure to glyphosate were reported: 1.2 (95% CI, 0.6–2.1; 27 exposed cases) for all lymphoid neoplasms combined, 1.0 (95% CI, 0.5–2.2; 12 exposed cases) for NHL, 0.6 (95% CI, 0.2–2.1; 4 exposed cases) for lymphoproliferative syndrome, 2.4 (95% CI, 0.8–7.3) for multiple myeloma, and 1.7 (95% CI, 0.6–5.0; 6 exposed cases) for Hodgkin lymphoma, after adjusting for age, and socioeconomic category.

•Cocco et al. (2013)⁶² performed a pooled analysis of case–control studies from six European countries to investigate the role of occupational exposure to specific groups of chemicals in the causation of lymphoma overall, B-cell lymphoma, and its most prevalent subtypes. A total of 2348 incident cases of lymphoma and 2462 controls were included in the study. Analyses were conducted for lymphoma and the most prevalent lymphoma subtypes and adjusted for age, sex, and education. A positive, non-significant association (odds ratio) of 3.1 (95% CI, 0.6–17.1) was reported for exposure to glyphosate and B-cell lymphoma.

I would note that the findings in the McDuffie et al. (2001)¹⁵; and Eriksson et al.¹⁸ studies is significant because their results are supported by the results reported for micronucleus formation studies in the bone marrow of mice by Rank et al. (1993)⁶³ where a single dose caused no effect while Bolognesi et al. (1997)³² and Manas et al. (2009)²⁷ reported that two daily doses of glyphosate did cause micronucleus formation in the bone marrow of mice in their studies. **This implies that level of exposure is an important consideration in the formation of NHL from exposure to glyphosate.**

Cohort Studies

The Agricultural Health Study (AHS)⁶⁴ is a large prospective study of cancer and other health outcomes in a cohort of licensed pesticide applicators and their spouses from Iowa and North Carolina. The AHS began in 1993 with the goal of answering important questions about how agricultural, lifestyle and genetic factors affect the health of farming populations. More than 89,000 farmers and their spouses in Iowa and North Carolina have participated in the study. It is the only cohort study to date to have published findings on exposure to glyphosate and the risk of cancer at many different sites. My summary of the 7 papers available evaluating cancer incidence associated with pesticide use in the AHS cohort follows:

- No risk estimates and no significant exposure-response associations with cancer of the prostate and exposure to glyphosate were reported by Alavanja et al (1996).⁶⁵

- DeRoos et al. (2005)^{66,67} evaluated associations between glyphosate exposure and the incidence of cancer at multiple sites in this cohort including lung, melanoma, multiple myeloma, and NHL, oral cavity, colon, rectum, pancreas, kidney, bladder, prostate, and leukemia. No significant exposure–response association with cancer at any of these sites was found.

- Flower et al.,⁶⁸ reported the results of the analyses of risk of childhood cancer associated with pesticide application by the parents of 17,357 children of Iowa pesticide applicators from the AHS cohort. For all the children of the pesticide applicators, the risk of cancer was increased for all childhood cancers combined, for all lymphomas combined, and for Hodgkin lymphoma, compared with the general population. A non-significant association (odds ratio) for use of glyphosate and risk of childhood cancer was reported to be 0.61 (95% CI, 0.32–1.16; 13 exposed cases) for maternal use and 0.84 (95% CI, 0.35–

2.34; 6 exposed cases) for paternal use.

- The incidence of cancer of the breast among farmers' wives in the AHS cohort, which included 30,454 women with no history of cancer of the breast before enrolment was reported by Engel et al.,⁶⁹. There was no difference in incidence of breast cancer for women who reported ever applying pesticides compared with the general population. A non-significant association (relative risk) for cancer of the breast was reported to be 0.9 (95% CI, 0.7–1.1; 82 cases) among women who had personally used glyphosate and a non-significant positive association (relative risk) of 1.3 (95% CI, 0.8–1.9; 109 cases) among women who never used pesticides but whose husband had used glyphosate.

- Lee et al.,⁷⁰ studied the relationship between exposure to agricultural pesticides and incidence of cancer of the colorectum in the AHS cohort. Non-significant positive associations (relative risks) with exposure to glyphosate was reported to be 1.2 (95% CI, 0.9–1.6) for cancers of the colorectum, and 1.6 (95% CI, 0.9–2.9) for cancers of the rectum. A non-positive association of 1.0 (95% CI, 0.7–1.5) was reported for cancers of the colon.

- Andreotti et al.,⁷¹ used a case–control analysis nested in the AHS cohort to study associations between the use of pesticides and cancer of the pancreas. For pancreatic cancer, a positive association (odds ratio) for ever- versus never-exposure to glyphosate was found but not significant at 1.1 (95% CI, 0.6–1.7; 55 exposed cases) and for highest category of level of intensity-weighted lifetime days was also found but not significant at 1.2 (95% CI, 0.6–2.6; 19 exposed cases).

- Dennis et al.,⁷² reported that exposure to glyphosate was not associated with cutaneous melanoma within the AHS cohort but did not report a risk estimate.

Meta-Analyses

- Schinasi & Leon⁷³ conducted a systematic review and meta-analysis of NHL and occupational exposure to agricultural pesticides, including glyphosate. The meta-analysis for glyphosate included six studies (McDuffie et al.,¹⁵ Hardell et al.,¹⁷ DeRoos et al.,^{67,11} Eriksson et al.,¹⁸ and Orsi et al.⁶¹) and yielded a significant positive association (meta risk-ratio) of 1.5 (95% CI, 1.1–2.0) for exposure to glyphosate and NHL.

- IARC⁷⁴ conducted an additional meta-analysis of NHL and occupational exposure to agricultural pesticides, including glyphosate using data from Schinasi & Leon⁷³ and

included the fully adjusted risk estimates from the studies published by Hardell et al.,¹⁷ and Eriksson et al.¹⁸ After considering the adjusted estimates of the two Swedish studies in the meta-analysis, the positive association (meta risk-ratio) was still significant at 1.3 (95% CI, 1.03–1.65).

•Chang and Delzell³ also conducted a systematic review and meta-analysis to examine the relationship between glyphosate exposure and risk of lymphohematopoietic cancer including NHL, Hodgkin lymphoma, multiple myeloma, and leukemia. Their analysis showed a positive association (meta-relative risks or meta-RRs) and was statistically significant for the association between any versus no use of glyphosate and risk of NHL (meta-RR=1.3, 95% confidence interval (CI)=1.0–1.6, based on six studies) and multiple myeloma (meta-RR =1.4, 95% CI=1.0–1.9; four studies). The authors conducted four meta-analyses for NHL, all reporting to have a significant positive association (meta-RR) of 1.3 or 1.4 with 95% CIs ranging from (1.0-1.6) to (1.0-1.8). The authors concluded “we found marginally significant positive meta-RRs for the association between glyphosate use and risk of NHL.”

Summary for Human Data

I have evaluated available epidemiology data. Based on my experience doing hazard assessments, I learned that epidemiologists consider case–control studies particularly valuable for determining the carcinogenicity of an agent because their design facilitates exposure assessment and reduces the potential for certain biases. My review of the literature finds that the two case-control studies from the United States and Canada, and the two case–control studies from Sweden indicated statistically significant positive associations between exposure to glyphosate and NHL. The Canadian study, McDuffie (2001)¹⁵, reported a positive association between glyphosate exposure and NHL for those case subjects with more than two days/year of exposure (odds ratio of 2.12(95%CI,1.20–3.73) when compared to those with less than two days exposure. Three studies reported excesses for NHL associated with exposure to glyphosate, after adjustment for other pesticides, De Roos (2003) reported a significant positive association (odds ratio) for a pooled US study¹¹ at 2.1 (95% CI, 1.1–4.0).; and the two Swedish studies (Hardell (2002)¹⁷, Eriksson (2008)¹⁸) reported significant positive associations of 3.04; 95% CI, 1.08–8.52

and 2.36 (95% CI, 1.04–5.37). The positive association from Hardell (2002)¹⁷ decreased to non-significance (1.85 (95% CI, 0.55–6.2)) when study area, and vital status were considered. Subtype-specific analyses in a Eriksson (2008)¹⁸ indicated positive associations for total NHL, as well as all subtypes, but this association was statistically significant only for the subgroup of lymphocytic lymphoma/chronic lymphocytic leukemia (odds ratio, 3.35; 95% CI, 1.42–7.89). A European study⁶² based on few cases also indicated an elevated risk (OR, 3.1; 95% CI, 0.6–17.1) for B-cell lymphoma. A French hospital-based case–control study⁶¹ did not find an association between exposure to glyphosate and NHL (OR, 1.0; 95% CI, 0.5–2.2) based on few exposed cases. For the evaluation of glyphosate, the Agricultural Health Study (AHS) is currently the only cohort study available providing information on its potential carcinogenicity and did not show an excess of NHL. There were three groups that did meta-analyses of the human data for an association between glyphosate use and NHL. Schinasi and Leon⁷³ reported a significant positive association (meta-RR) of 1.5 (95% CI, 1.1–2.0). The IARC study⁷⁴ showed a positive association (meta-RR) of 1.3 (95% CI, 1.03–1.65). Chang and Delzel³ provided four separate meta-analyses, all of which are reported as having a significant association (meta-RR) of either 1.3 or 1.4 with CIs ranging from (1.0–1.6) to (1.0–1.8). When the data across all epidemiological studies are combined, results indicate a positive association between glyphosate exposure and NHL in humans.

Interpreting the epidemiology findings requires one to properly weight studies according to quality rather than simply count the number of positives and negatives. The pooled case–control analysis from the USA¹¹ contained 650 cases of NHL. It follows that the case-control studies provide a stronger assessment of the potential carcinogenicity of glyphosate. The case-control studies in the US¹¹, Canada¹⁵ and Sweden^{17,18} indicate a significant positive association for NHL with exposure to glyphosate. This positive association was also observed in the studies that adjusted for other pesticides. The AHS cohort did not show an excess of NHL; however it reports on only 92 NHL cases in the unadjusted analysis.⁶⁴ The three meta-analyses I reviewed are good examples of objective evaluations and show a consistent positive association between glyphosate and NHL. Drawing on the Bradford-Hill criteria⁷⁵ for causality, I would state that the observations are consistent (relative risks and meta analyses are positive for the case control studies), significant, not specific, temporally observed, shows a biological gradient, and is coherent

with the animal evidence (discussed below). Using my stated criteria, I conclude there is “Limited” evidence for the carcinogenicity of glyphosate in humans, because a positive association has been observed between exposure to glyphosate and NHL, and a causal interpretation is creditable but alternative explanations such as chance, bias or confounding could not be completely ruled out.

Hazard Assessment of the Experimental Animal Data for Glyphosate and/or Glyphosate-Based Formulations

Before discussing the experimental animal data for glyphosate and/or glyphosate-based formulations, I will define what is involved in a cancer bioassay in experimental animals. The basic cancer bioassay design has remained relatively constant for more than 40 years and consists of groups of 50 male and female mice and rats in each dose and control group. Treatment traditionally lasts for 24 months and commences when the animals are 6–8 weeks of age. Early bioassay studies involved two treatment groups plus a control group. The first treatment group was a high dose, referred to as a maximally tolerated dose (MTD), and the second treatment group was half that dose. More recent studies typically include three (and sometimes up to five) treatment groups plus the control group.

In the bioassays, I reviewed the nature and extent of impurities or contaminants, the animal species, strain, sex, numbers per group, age at start of treatment, route of exposure, dose levels, duration of exposure, survival and information on tumors. With regard to the tumors, I evaluated the incidence, latency, severity or multiplicity of neoplasms or preneoplastic lesions. Studies in experimental animals that I determined to be inadequate for evaluation (e.g. too short a duration, too few animals, poor survival) can be found at the end of my reference list.

Cancer Bioassays in Mice

•Knezevich and Hogan⁷⁶ (1983) were the authors of a report submitted to the Environmental Protection Agency (EPA)⁷⁷ by Monsanto in support of the registration of glyphosate as an herbicide. This report was also discussed in the paper by Greim⁷⁸ (referred to as Study 10). For 24 months, groups of 50 male and 50 female CD-1 mice received diets containing glyphosate (purity, 99.7%) at a concentration of 0, 1000, 5000,

or 30,000 ppm, ad libitum. The study observed no treatment-related effect on body weight in male and female mice at the lowest or intermediate dose, but a slight reduction in body weight in the male and female mice at the highest dose compared with controls. Survival in all dose groups was similar to controls. (It does not appear that a MTD was reached). There was a positive trend⁷⁹ ($p = 0.016$, trend test) in the incidence of renal tubule adenoma in dosed male mice: 0/49, 0/49, 1/50 (2%), 3/50 (6%). Renal tubule adenoma is a rare tumor in CD-1 mice. Historical control data from 14 studies conducted between 1977 and 1981 at the testing laboratory indicated that the mouse renal tumors ranged from 0 to 3% and the incidence in the current study (3/50; 6%) exceeded the upper limit of the historical control range by a factor of two. The rarity of this tumor in CD-1 mice is documented in a publication by Chandra and Frith⁸⁰ that reports only 1 out of 725 [0.14%] CD-1 male mice in their large historical database had developed renal cell tumors (one carcinoma). No tumors of the kidney were observed in the female mice. No other tumor sites were identified.

A re-evaluation of the original renal section was conducted by a Monsanto consulting pathologist who reported a small renal tubule adenoma in one control male mouse, which was not diagnosed as such in the original pathology report.⁸¹ This finding was contrary to the initial findings of Bio/dynamics lab, the lab commissioned to complete this report. Following Monsanto's submission of the consulting pathologist's report, the EPA reported there was no difference in diagnoses between his and other pathologists' diagnoses with respect to kidney tumors in mid- and high-dose groups (i.e. 0/49, 0/49, 1/50 (2%), 3/50 (6%)). The EPA pathologist also indicated in his report⁷⁹ this data also shows a positive trend ($p = 0.016$, trend test) in the incidence of renal tubule adenoma in the dosed male mice. Regarding the questionable male control kidney, it was his opinion that the presence of a tumor cannot definitely be established. Nonetheless, the EPA requested additional renal sections be cut and evaluated from all male mice in the control and treated groups; this additional review found no additional tumors.⁸¹ The EPA also requested that a pathology working group (PWG) be convened to evaluate the tumors of the kidney observed in male mice treated with glyphosate, including the additional renal sections.⁸² Monsanto sponsored a PWG that reported the incidence of adenoma of the renal tubule was 1/49 (2%), 0/49, 0/50, 1/50 (2%)(not statistically significant); the incidence of carcinoma of the renal tubule was 0/49, 0/49, 1/50 (2%),

2/50 (4%) (which gives a significant $p = 0.037$, trend test for carcinoma); and the incidence of adenoma or carcinoma (combined) of the renal tubule was 1/49 (2%), 0/49, 1/50 (2%), 3/50 (6%) (which gives a significant $p = 0.034$, trend test for combined). The PWG did not discuss their finding of an adenoma in the control male mice or address the previous opinion that the presence of a tumor in the control male mice cannot definitely be established and concluded the kidney tumors were not compound related.⁸³ It is important to note that the renal tumor identified in the controls by the PWG after re-evaluation of the original slides was not seen in the re-sectioned kidney slides. **My conclusion of the results discussed above is that there was a significant increase in the incidence of these rare kidney tumors in the CD-1 mouse, with a dose-related trend, which is caused by glyphosate.** For the purpose of this hazard identification the increase the incidence of carcinoma of the renal tubule and the incidence of adenoma or carcinoma (combined) of the renal tubule in male mice is due to treatment with glyphosate that caused a significant, dose related increase of these rare tumors in male CD-1 mice.

•Atkinson et al.⁸⁴ (1993) were the authors of a report submitted to the EPA in support of the re-registration of glyphosate as an herbicide. This study was also discussed in the paper by Greim⁷⁸ (Study 11). Groups of 50 male and 50 female CD-1 mice were given diets containing glyphosate (purity, 98.6%) at a concentration that was adjusted weekly for the first 13 weeks and every 4 weeks thereafter to give doses of 0, 100, 300, or 1000 mg/kg bw, ad libitum, for 104 weeks. There was no treatment-related effect on body weight or survival in any of the dosed groups indicating a maximum tolerated dose was not achieved. The EPA reported⁷⁷ a statistically significant increase in the incidence of hemangiosarcoma (blood vessel tumor) in males – 0/47, 0/45, 0/50, 4/45 (9%) ($p < 0.01$, trend test), and non-significant increase in females – 0/50, 2/50 (4%), 0/50, 1/50 (2%). The EPA pointed out that the incidence in the high dose males was near the upper limit (0-8%) for the performing laboratory. However, if one looks at excerpts from the full report,⁸⁴ Table 15 (page 97) indicates that as few as 2 animals per dose group were examined histologically for this tumor. This would lead one to consider that the incidence of this tumor could have been higher in this study as more of these tumors could have been found if all 50 animals per dose group were examined. There was also reported a non-significant increase in the incidence of histiocytic sarcoma in the lymphoreticular/haemopoietic tissue in males – 0/50, 2/50 (4%), 0/50, 2/50 (4%), and

in females – 0/50, 3/50 (6%), 3/50 (6%), 1/50 (2%). The EPA stated⁷⁷ that for their risk analysis, the increase in hemangiosarcomas in male mice was not considered to be treatment-related. For the purpose of this hazard identification, I determined the increased incidence of hemangiosarcomas in male mice is due to the treatment with glyphosate that caused a significant dose related increase in the incidence of hemangiosarcoma in male CD-1 mice. This association may have been stronger if all the animals in this study had been examined histologically for this tumor.

•Greim⁷⁸ (Study 12, Sugimoto, K.) reported on a study submitted by Arysta Life Sciences to the EPA in support of the re-registration of glyphosate as an herbicide. Groups of ICR-CD-1 mice (50/sex/group received diets containing glyphosate (94.6–97.6% pure) at 0, 1600, 8000 or 40,000 ppm for 18 months. Parameters evaluated included clinical signs, body weight, food consumption, hematology, clinical chemistry, and urinalysis, organ weights, gross necropsy and histopathological examination. The EPA reported⁷⁷ no adverse effects on survival were observed in either sex across the doses tested and there were no statistically significant increases in any tumor type in this study based on details provided by Greim⁷⁸. A review of the tumor tables for this study (Sugimoto⁸⁵) shows that there was a significant trend for the development of hemangiosarcomas in male mice (0/50; 0/50; 0/50; 2/50 (4%)) with a p-value for trend of 0.008, Chi-Square test; a significant trend for the development of malignant lymphomas in male mice (2/50 (4%); 2/50 (4%); 0/50; 6/50 (12%)) with a p-value for trend of 0.008, Chi-Square test; and a significant trend for the development of renal adenomas (0/50; 0/50; 0/50; 2/50 (4%)) with a p-value for trend of 0.008, Chi-Square test seen in male mice. The EPA also reported⁸⁶ that hemangiosarcomas in female mice were found to occur with a statistically significant trend in this study (0/50; 0/50; 2/50, (4%); 5/50, (10%) $p=0.002$, Trend test), and the tumor incidence in the high-dose female mice was statistically significant with $p=0.028$ as compared to concurrent controls. I also reviewed the Tier II Summaries for Glyphosate Carcinogenicity Studies from Greim, et al.⁸⁷ for Study 12, Sugimoto, which showed a reported statistically significant increase in malignant lymphoma in high dose male mice – 0/26, 0/34, 1/27(4%), 5/29(17%) ($p<0.05$ Fisher's exact test); however I could not resolve the difference in the tumor incidence between the Greim Tier II Summary⁸⁷, the published Greim et al, 2015⁷⁸ and the Sugimoto⁸⁵ tumor tables. These appear to be low response rates but this is only an 18-month study where low rates of

tumors are not unusual. For the purpose of this hazard identification there was an increased incidence of malignant and/or a combination of malignant and benign tumors, at multiple tissue sites in male and female CD-1 mice in this study. The significant increase in malignant lymphoma in high dose male mice, and the significant trend in the development of hemangiosarcomas, malignant lymphomas, and renal adenomas in male mice is due to treatment with glyphosate that caused these cancers in male CD-1 mice. The significant trend in the development of hemangiosarcomas in female mice is also related to treatment with glyphosate that caused this cancer in female CD-1 mice.

•Greim⁷⁸ (Study 14, Wood, et al. 2009b) reported on a study submitted by Nufarm to the EPA in support of the re-registration of glyphosate as an herbicide. Groups of 51 male and 51 female CD-1 mice were given diets containing glyphosate (purity, 94.6–97.6%) at a concentration of 0, 500, 1500, or 5000 ppm for 18 months. Parameters evaluated included clinical signs, body weight, food consumption, organ weights, gross necropsy and histopathological examination. There was no treatment-related effect on survival. In male mice at the high dose there was a significant increase in the incidence of malignant lymphomas (0/51, 1/50(10%), 2/51(4%), 5/51(10%) $p < 0.05$, pair-wise comparison, $p < 0.01$ for trend) and a significant increase in the trend of formation of adenocarcinomas of the lung (5/51(10%), 5/51(10%), 7/51(14%), 11/51(22%) $p < 0.01$ for trend⁸⁶). For the purpose of this hazard identification, I determined the formation of malignant lymphomas and the formation of adenocarcinomas of the lung in male mice in this study is due to treatment with glyphosate that caused a significant increase in the incidence of malignant lymphoma in high dose male CD-1 mice and an increase in the trend of formation of the adenocarcinomas of the lung and malignant lymphomas in male CD-1 mice.

•Greim⁷⁸ (Study 13, Kumar) reported on a study submitted by Feinchemie Schwebda to the EPA in support of the re-registration of glyphosate as an herbicide. Groups of 50 male and 50 female Swiss albino mice [age at start not reported] were given diets containing glyphosate (purity >95%) at a concentration of 0, 100, 1000, or 10,000 ppm for 18 months. There were no treatment-related effects on clinical signs, behavior, body weight, body weight gain, food consumption, and differential white blood cell counts in both sexes. There was a slightly higher mortality rate observed in the high dose groups. There was a significant increase in malignant lymphoma reported in high dose male mice

(10/50, 20%; 15/50, 30%; 16/50, 32%; 19/50, 38%, $p < 0.05$ pair wise) and female mice (18/50, 36%; 20/50, 40%; 19/50, 38%; 25/50, 50%, $p < 0.05$ pair wise). There was also a significant increased trend (one-sided p -value for trend = 0.05) for the formation of this tumor in males. The incidence of malignant lymphoma in the high dose male was double the historical rate, reported to be 18%⁸⁷ for males, and for high dose female mice the incidence was well above the historical rate of 41%⁸⁷. There was also a significant increased trend in the incidence of kidney renal cell adenomas reported⁸⁸ in males (0/50; 0/26; 1/26 (4%); 2/50 (4%); one-sided p -value for trend $p = 0.04$). I would note that the EPA stated⁷⁷ this study was not included in their review due to the report by Greim (2015)⁷⁸ that there was possibly a viral infection within the colony, which confounded the interpretation of the study findings. EPA also stated although the incidences in this study were within or near the normal variation of background occurrence. It is not clear whether or not the viral component may have contributed to incidence value reported or the lower survival seen at the high dose in the study.⁸⁹ An internal Monsanto email among the authors of Greim would indicate there was no viral infection in the mouse colony during this study. Further, Greim⁷⁸ (table 18, p. 201) considers this study GLP and OECD compliant. For the purpose of this hazard identification, I determined formation of malignant lymphoma in the male and female mice and the renal cell adenomas in males in this study is due to treatment with glyphosate that caused a significant increase in the incidence of malignant lymphoma in high dose male and female Swiss albino mice and renal cell adenomas in male Swiss albino mice.

Cancer Bioassays in Rats

•Greim⁷⁸ reported on a Bio/dynamics study (Study 1, Lankas, et al.) submitted by Monsanto to the EPA in support of the registration of glyphosate as an herbicide. Groups of 50 male and 50 female Sprague-Dawley rats were fed diets containing glyphosate (98.7%, pure) at concentrations of 0, 30, 100 or 300 ppm for 26 months. These concentrations were adjusted during the course of the study so that actual doses of 0, 3, 10, and 31 mg/kg/day in males and 0, 3, 11, and 34 mg/kg/day in females were maintained. There were no treatment-related effects on body weight or survival at any dose level. An MTD was not achieved. There was a significant increase reported in the incidences of interstitial cell tumors in the testes of male rats: controls 0/50, 0%; low dose

3/5, 6%; mid dose 1/50, 2%; high dose 6/50; 12%; $p=0.013$ by pairwise comparison⁷⁷. The incidence of interstitial cell tumors in the testes in the high dose animals in this study is almost twice that seen in the range of this tumor (3.4% to 6.7%) in control animals (historical controls) from 5 contemporary studies⁸⁷. There was also a significant increase in the incidence of pancreatic islet cell adenoma reported in males at the low dose: controls, 0/50; low dose 5/49, 10% ($p < 0.05$ Fisher exact test); mid dose 2/50, 4%; high dose 2/50, 4%. For the purpose of this hazard identification, I determined the increase in the incidence of interstitial cell tumors in the testes and pancreatic cell tumors in male rats is due to the treatment with glyphosate that caused a significant increase in the incidence of interstitial cell tumors in the testes and pancreatic islet cell tumors in male Sprague-Dawley rats.

• Greim⁷⁸ reported on a study (Study 2, Stout, et al.) submitted by Monsanto to the EPA in support of the registration of glyphosate as an herbicide. Groups of 60 male and 60 female Sprague-Dawley rats were given diets containing glyphosate (technical grade; purity, 96.5%) at a concentration of 0 ppm, 2000 ppm, 8000 ppm, or 20,000 ppm, ad libitum, for 24 months. No compound-related effect on survival was observed. There was no statistically significant decreases in body-weight gain in male rats. The study reported significant decreases in body-weight gain in females at the highest dose, beginning on day 51. There was a statistically significant increase in the incidence of pancreatic islet cell adenoma in males at the lowest dose compared with controls: control 1/58, 2%; low dose 8/57, 14% ($p \leq 0.05$ Fisher exact test); mid dose 5/60, 8%; high dose 7/59, 12%. The EPA⁷⁷ did additional analysis of this data for pancreatic islet cell adenoma by excluding rats that died or were killed before week 55 and then using static analyses (Cochran–Armitage trend test and Fisher exact test) that gave a statistically significant higher incidence of these tumors in males at the lowest and highest doses compared with controls: control 1/43, 2%; low dose 8/45, 18% ($p = 0.018$; pairwise test); mid dose 5/49, 10%; high dose 7/48, 15% ($p = 0.042$; pairwise test). The incidence of these adenomas in the low (18%) and high (15%) dose males was almost twice that seen in historical controls. The range for historical controls for pancreatic islet cell adenoma reported in males at this laboratory was 1.8–8.5%⁷⁷. One should note that there was no statistically significant positive trend in the incidence of these tumors, and no apparent progression to carcinoma. There was also a statistically significant positive trend ($p = 0.016$) in the

incidence of hepatocellular adenoma observed in male rats⁸⁶ and a statistically significant positive trend of thyroid follicular cell adenomas ($p = 0.031$) and thyroid follicular cell adenomas and carcinomas combined ($p=0.033$) observed in female rats⁸⁶ reported in this study. For the purpose of this hazard identification, I determined that the increase in the incidence of pancreatic islet cell adenoma in male rats is due to the treatment with glyphosate that caused a significant positive increase in the incidence of pancreatic islet cell adenomas of male Sprague-Dawley rats. Glyphosate also caused a significant increase in the trend for formation of hepatocellular adenomas in male Sprague-Dawley rats and of thyroid follicular cell adenomas and follicular cell adenomas and carcinomas combined in female Sprague-Dawley rats.

•Greim⁷⁸ reported on a study (Study 3, Atkinson, et al.) submitted by Cheminova to the EPA in support of the registration of glyphosate as an herbicide. Groups of 50 male and 50 female Sprague-Dawley rats were given diets containing glyphosate, purity, 98.7–98.9%, at a concentration that were adjusted to provide doses of 0, 10, 100, 300, or 1,000 mg/kg bw/day, ad libitum, for 104 weeks. Decreased body-weight gain was observed in males and females at the highest dose. There was no significant decrease in survival reported at any dose level. Neoplasms were noted in control and treated groups, but dose-responses were not evident, and no statistically significant increases versus controls were noted for any tumor type. Additionally, EPA's evaluation⁸⁶ of this study indicated there were no treatment-related increases in the occurrence of any tumor type in this study.

•Greim⁷⁸ reported on a study (Study 7, Brammer) submitted by Syngenta to the EPA in support of the re-registration of glyphosate as an herbicide. Groups of 52 male and 52 female Wistar rats received diets containing 0, 2,000, 6,000, and 20,000 ppm glyphosate (97.6% pure), ad libitum, for 24 months. Survival in the high dose group males was significantly better than the other dose groups throughout the study while survival in the females was similar across all dose groups. The bodyweights of the high dose males and females were statistically significantly lower than controls throughout the study. The study's author reported no significant increase in tumor incidence in any of the treated groups. The EPA's evaluation⁷⁷ of this study indicated there was a significant increase in the incidence of hepatocellular adenomas in male rats at the high dose when compared to controls (control 0/52, 0%; low dose 2/52, 4%; mid dose 0/52, 0%; high dose 5/52, 10%, $p=0.03$). There was also a significant trend ($p=0.008$) in the formation of this tumor in

male rats. The EPA goes on to state the incidences observed were within the range (0–11.5%) of historical controls for this strain of rats in 26 studies conducted during the relevant time period (1984–2003) at the testing laboratory indicating this increase was not considered to be related to treatment with glyphosate. For the purpose of this hazard identification, I determined the increase in the formation of hepatocellular adenomas in male Wistar rats could not be attributed to exposure to glyphosate in this study despite the fact that there was an observation of increased incidence of hepatocellular adenomas in male rats.

- Greim⁷⁸ reported on a study (Study 4, Suresh) submitted by Feinchemie Schwebda to the EPA in support of the registration of glyphosate as an herbicide. Groups of 50 male and 50 female Wistar rats received diets containing 0, 100, 1,000, and 10,000 ppm glyphosate (97.6% pure), ad libitum, for 24 months. There were no treatment-related deaths or clinical signs in any of the dose-groups and there were no treatment related effects on body weight gain or food consumption noted. This suggests that the MTD was not reached, and this study is inadequate for the evaluation of the carcinogenicity of glyphosate.

- Greim⁷⁸ reported on a study (Study 6, Enomoto) submitted by Arista Life Sciences to the EPA in support of the registration of glyphosate as an herbicide. Groups of 50 male and 50 female Sprague-Dawley rats received diets containing 0, 3,000, 10,000, or 30,000 ppm glyphosate (94.6–97.6% pure) for 24 months. Decreases in body weight were observed in both sexes in the mid and high dose group along with a lower food consumption. Survival in the high dose males was lower than controls while there was no compound-related effect on survival in any other dose group. There were no statistically significant increases in any tumor type reported for this study.

- Greim⁸² reported on a study (Study 8, Wood 2009a) submitted by Nufarm to the EPA in support of the registration of glyphosate as an herbicide. Groups of 51 male and 51 female Wistar rats received diets containing 0, 3,000, 10,000, or 15,000 ppm glyphosate (95.7% pure) for 24 months, the highest dose level was progressively increased to 24000 ppm by week 40. There were no treatment-related deaths or clinical signs in any of the dose-groups. No significant treatment-related effects on mortality were observed during the study. This suggests that the MTD was not reached, and this study is inadequate for the evaluation of the carcinogenicity of glyphosate.

•Chruscielska et al.⁹⁰ gave groups of 55 male and 55 female Wistar rats drinking-water containing an ammonium salt of glyphosate (purity not given) that was used to make drinking water solutions of 0, 300, 900, and 2700 mg/L, for 24 months. The authors reported that survival and body-weight gain were similar in treated and control animals and that no significant increase in tumor incidence was observed in any of the treated groups. There was limited information provided on dosing regimen, histopathological examination method, and tumor incidences that makes this study inadequate for the purpose of this hazard assessment.

Summary for Experimental Animal Data

I reviewed a total of five dose feed bioassays of glyphosate in mice. Four of these studies (Study 12 and Study 14 in Greim⁷⁸, Knezevich and Hogan (1983)⁷⁶, and Atkinson⁸⁴) were in male and female CD-1 mice, and one study^{78(Study 13)} was in male and female Swiss albino mice. Glyphosate caused a significant increase in the incidence of adenoma or carcinoma (combined) and a significant positive trend for the formation of adenoma or carcinoma (combined) of the renal tubule in male CD-1 mice in one study⁷⁶, and a significant positive trend for the formation of adenomas of the renal tubule in male CD-1 mice in another study^{78(Study 12)}. Glyphosate also caused a significant increase in the incidence of renal cell adenomas in male Swiss albino mice^{78(Study 13)}. Adenoma and carcinoma of the renal tubule constitutes a morphological continuum in the development and progression of renal neoplasia in mice^{91,92}. It is important to note that renal tubule carcinoma is a very rare tumor in CD1 mice⁸⁰ and that this tumor was caused by exposure to glyphosate in two different strains of mice (CD-1 and Swiss). Glyphosate caused a significant increase in the incidence of malignant lymphoma in male CD-1 mice in two studies^{78(Study 12, Study 14)} and in male and female Swiss albino mice in another study^{78(Study 12)}. Glyphosate also caused a significant positive trend for the formation of malignant lymphoma in one of these studies in male CD-1 mice^{78(Study 12)} and caused a significant positive trend for the formation of hemangiosarcomas in 2 separate studies in male CD-1 mice^{78(Study 12),84}. There was also a significant positive trend for the formation of adenocarcinomas of the lung in male CD-1 mice in one study^{78(Study 14)} and hemangiosarcomas in female CD-1 mice in another study^{82(Study 12)}.

I reviewed a total of 7 dosed feed and 2 drinking water bioassays of glyphosate in rats. Four of the feed studies and one drinking water study were in male and female Sprague-Dawley rats and three feed studies and one drinking water study were in male and female Wistar rats. Glyphosate caused a significant increase in the incidence of pancreatic islet cell adenoma in two feeding studies in male Sprague-Dawley rats^{78(Study 1 and Study 2)}. Glyphosate caused a significant increase in the incidence of thyroid tumors in male Sprague-Dawley rats in one feeding study^{78(Study 1)} and a significant positive trend for the formation of thyroid tumors in female Sprague-Dawley rats in another feeding study^{78(Study 2)}. Glyphosate caused a significant increase in the incidence of interstitial cell tumors in the testes of male Sprague-Dawley rats in one feeding study and a significant positive trend for the formation of hepatocellular adenomas in male Sprague-Dawley rats in another feeding study^{78(Study 1)}.

To state my findings more concisely, I determined that in CD-1 mice, glyphosate exposure causes kidney tumors in males in two separate studies^{76,78(Study 12)}, hemangiosarcomas in males in two separate studies,^{78(Study 12),84} malignant lymphoma in males in two separate studies^{78(Study 12, Study 14)}, adenocarcinomas of the lung in males in one study^{78(Study 14)}, and hemangiosarcomas in females in one study^{78(Study 12)}. In one study^{78(Study 13)} in Swiss albino mice, exposure to glyphosate causes malignant lymphoma in males and females and kidney tumors in males.

I also determined that in Sprague-Dawley rats, glyphosate exposure causes pancreatic cell tumors in males in one study^{78(Study 2)}, interstitial cell tumors in the testes in males in one study^{78(Study 1)}, hepatocellular adenomas in males in two studies^{78(Study 2, Study 7)}, and thyroid follicular cell tumors in females in one study^{78(Study 2)}.

Considering all data from the mice and rat studies I reviewed, there is "Sufficient" evidence that shows glyphosate is carcinogenic in experimental animals causing kidney tumors, hemangiosarcomas, malignant lymphoma, adenocarcinomas of the lung, and hemangiomas in mice and pancreatic cell tumors, interstitial cell tumors in the testes, hepatocellular adenomas, and thyroid follicular cell tumors in rats. This statement is based on my stated criteria of a causal relationship between exposure to glyphosate and an increased incidence of malignant and/or a combination of malignant and benign tumors, in multiple species, at multiple tissue sites, from multiple studies, and to an unusual degree with regard to incidence, site, or type of tumor.

Hazard Assessment of the Mechanistic and Other Data for Glyphosate and Glyphosate-Based Formulations

Data on the absorption of glyphosate via intake of food and water in humans could not be found in the published literature. Glyphosate has been found in the urine of agricultural workers. In a study by Acquavella⁷, 60% of farmers had detectable levels of glyphosate in 24-hour composite urine samples taken on the day they had applied a glyphosate-based formulation. Wearing protective gear such as rubber gloves reduced the concentrations of glyphosate in the urine. This implies that dermal absorption is a relevant route of exposure. Curwin⁸ demonstrated that glyphosate is also present in the urine of non-farm families. No data in humans on the distribution of glyphosate in systemic tissues other than blood were found in the available published literature. In cases of accidental or deliberate intoxication involving ingestion of glyphosate-based formulations, glyphosate was measured in blood.

Strong evidence indicates that glyphosate is genotoxic. As noted in Monograph 112, studies in human cells^{27,31,32}, mammalian model systems^{27,32,33}, and in non-mammalian organisms^{35,37} have given positive results. The end-points evaluated in these studies included biomarkers of DNA adducts and various types of chromosomal damage. Tests in bacterial assays gave consistently negative results.

The evidence for genotoxicity caused by glyphosate-based formulations is also strong. As noted in Monograph 112, three studies^{39,93,94} reported examining genotoxic end-points in community residents exposed to glyphosate-based formulations and two of these studies reported positive associations. One study³⁹ looked at micronucleus formation in circulating blood cells before and after aerial spraying with glyphosate-based formulations to determine chromosomal damage in exposed individuals. This study revealed a significant increase in micronucleus formation after exposure in three out of four different geographical areas. Additional positive evidence came from in vitro studies with positive results in human cells^{32,45}, in vivo^{27,32} and in vitro⁹⁵ studies in mammalian systems, and studies in non-mammalian organisms^{35,96} such as fish. Biomarkers of DNA adducts and different types of chromosomal damage were examined in these studies. The pattern of tissue specificity of genotoxicity end-points observed with glyphosate-based

formulations is similar to that observed with glyphosate. Tests of glyphosate-based formulations in bacterial assays gave generally negative results.

There is strong evidence that glyphosate and glyphosate-based formulations induce oxidative stress. As noted in Monograph 112, evidence of oxidative stress comes from in vitro studies in human cells^{97,98} and in many in vivo studies^{32,42}, examining rodent tissues. Studies of oxidative stress and glyphosate in non-human mammalian experimental systems were conducted in rats and mice, and examined a range of exposure durations, doses, preparations (glyphosate and glyphosate-based formulations), administration routes and tissues. In these studies glyphosate caused free radicals and oxidative stress in mouse and rat tissues through alteration of antioxidant enzyme activity, depletion of glutathione, and increases in lipid peroxidation. In at least one of the studies in human cells the oxidative stress caused by glyphosate was ameliorated by co-administration of antioxidants⁴⁰. Similar findings of oxidative stress have been reported in fish and other aquatic species providing additional evidence for glyphosate-induced oxidative stress⁹⁹. Molecular epidemiology studies^{100,101} have documented that oxidative stress is a pathway to the formation of NHL in humans. Further, the in vitro studies in humans cells and in vivo and in vitro studies in rodents provides evidence that exposure to glyphosate causes oxidative stress. Logically it follows that there is a positive association between oxidative stress caused by glyphosate and glyphosate-based formulations and NHL observed in humans exposed to glyphosate-based formulations and that a causal interpretation is credible.


Hazard Assessment Conclusion

Based on the significant positive association observed in the studies discussed above, I conclude that there is evidence that glyphosate and glyphosate-based formulations are carcinogenic in humans. First, the human study data supports a positive association between exposure to glyphosate and glyphosate-based formulations and the development of NHL. Second, all the data from the animal bioassay studies provide evidence that glyphosate is carcinogenic in experimental animals. Third, the mechanistic data show that glyphosate and glyphosate-based formulations cause genotoxicity and oxidative stress in humans and animals. Therefore, I conclude to a reasonable degree of

scientific certainty that glyphosate and glyphosate-based formulations are probable human carcinogens. I also conclude to a reasonable degree of scientific certainty that glyphosate and glyphosate-based formulations cause NHL in humans.

Compensation and Testimony

My billing rate is \$400/hr plus travel fees and expenses. I have not testified in any case in the last four years.



Charles W. Jameson, Ph.D.

References

- ¹ Williams et al. (2016) A review of the carcinogenic potential of glyphosate by four independent expert panels and comparison to the IARC assessment. *Critical Reviews in Toxicology*, VOL. 46, NO. S1, 3–20
- ² Williams et al. (2016) Glyphosate rodent carcinogenicity bioassay expert panel review. *Critical Reviews in Toxicology*, VOL. 46, NO. S1, 44–55
- ³ Chang ET, and Delzell, E (2016) Systematic review and meta-analysis of glyphosate exposure and risk of lymphohematopoietic cancers. *J Environ Sci Health B* ;51(6):402-34.
- ⁴ Solomon KR, (2016) Glyphosate in the general population and in applicators: a critical review of studies on exposures *Critical Reviews in Toxicology*. VOL. 46, NO. S1, 21–27
- ⁵ See, IARC Preamble, <http://monographs.iarc.fr/ENG/Preamble/index.php> (2015)
- ⁶http://www.prweb.com/releases/glyphosate_agrochemical/technical_glyphosate/prweb8857231.htm
- ⁷ Acquavella JF, Alexander BH, Mandel JS, Gustin C, Baker B, Chapman P et al. (2004). Glyphosate biomonitoring for farmers and their families: results from the Farm Family Exposure Study. *Environ Health Perspect*, 112(3):321–6.
- ⁸ Curwin BD, Hein MJ, Sanderson WT, Striley C, Heederik D, Kromhout H et al. (2007). Urinary pesticide concentrations among children, mothers and fathers living in farm and non-farm households in Iowa. *Ann Occup Hyg*, 51(1):53–65.
- ⁹ MLHB (2013). Determination of glyphosate residues in human urine samples from 18 European countries. Bremen: Medical Laboratory of Bremen. Available from: https://www.foeeurope.org/sites/default/files/glyphosate_studyresults_june12.pdf,
- ¹⁰ Varona M, Henao GL, Díaz S, Lancheros A, Murcia A, Rodríguez N et al. (2009). Evaluación de los efectos del glifosato y otros plaguicidas en la salud humana en zonas objeto del programa de erradicación de cultivos ilícitos. [Effects of aerial applications of the herbicide glyphosate and insecticides on human health] *Biomedica*, 29(3):456–75.
- ¹¹ De Roos AJ, Zahm SH, Cantor KP, Weisenburger DD, Holmes FF, Burmeister LF et al. (2003). Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men. *Occup Environ Med*, 60(9):E11
- ¹² Zahm SH, Weisenburger DD, Babbitt PA, Saal RC, Vaught JB, Cantor KP et al. (1990). A case-control study of non-Hodgkin's lymphoma and the herbicide 2,4-dichlorophenoxyacetic acid (2,4-D) in eastern Nebraska. *Epidemiology*, 1(5):349–56.
- ¹³ Hoar SK, Blair A, Holmes FF, Boysen CD, Robel RJ, Hoover R et al. (1986). Agricultural herbicide use and risk of lymphoma and soft-tissue sarcoma. *JAMA*, 256(9):1141–7.

- ¹⁴ Cantor KP, Blair A, Everett G, Gibson R, Burmeister LF, Brown LM et al. (1992). Pesticides and other agricultural risk factors for non-Hodgkin's lymphoma among men in Iowa and Minnesota. *Cancer Res*, 52(9):2447-55.
- ¹⁵ McDuffie HH, Pahwa P, McLaughlin JR, Spinelli JJ, Fincham S, Dosman JA et al. (2001). Non-Hodgkin's lymphoma and specific pesticide exposures in men: cross-Canada study of pesticides and health. *Cancer Epidemiol Biomarkers Prev*, 10(11):1155-63
- ¹⁶ Kachuri L, Demers PA, Blair A, Spinelli JJ, Pahwa M, McLaughlin JR et al. (2013). Multiple pesticide exposures and the risk of multiple myeloma in Canadian men. *Int J Cancer*, 133(8):1846-58
- ¹⁷ Hardell L, Eriksson M, Nordstrom M (2002). Exposure to pesticides as risk factor for non-Hodgkin's lymphoma and hairy cell leukemia: pooled analysis of two Swedish case-control studies. *Leuk Lymphoma*, 43(5):1043-9
- ¹⁸ Eriksson M, Hardell L, Carlberg M, Akerman M (2008). Pesticide exposure as risk factor for non-Hodgkin lymphoma including histopathological subgroup analysis. *Int J Cancer*, 123(7):1657-63.
- ¹⁹ EPA (1985a). Glyphosate; EPA Reg.#: 524-308; Mouse oncogenicity study. Document No. 004370. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/103601/103601-183.pdf>, accessed 10 March 2015.
- ²⁰ EPA (1985b). EPA Reg.#: 524-308; Roundup; glyphosate; pathology report on additional kidney sections. Document No. 004855. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/clearedreviews/reviews/103601/103601-206.pdf>.
- ²¹ EPA (1986). Glyphosate; EPA Registration No. 524-308; Roundup; additional histopathological evaluations of kidneys in the chronic feeding study of glyphosate in mice. Document No. 005590. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/>
- ²² EPA(1991a). Second peer review of glyphosate. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/103601/103601-265.pdf>
- ²³ JMPR (2006). Glyphosate. In: Joint FAO/WHO Meeting on Pesticide Residues. Pesticide residues in food - 2004: toxicological evaluations. Report No. WHO/PCS/06.1. Geneva: World Health Organization; pp. 95-169. Available from: http://whqlibdoc.who.int/publications/2006/9241665203_eng.pdf?ua=1
- ²⁴ EPA (1991b). Glyphosate; 2-year combined chronic toxicity/carcinogenicity study in Sprague-Dawley rats - List A pesticide for reregistration. Document No. 008390.

Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/103601/103601-263.pdf>, see also <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/103601/103601-268.pdf>

²⁵ EPA (1991c). Peer review on glyphosate. Document No. 008527. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency.

²⁶ EPA (1991d). Glyphosate – EPA registration No. 524-308 – 2-year chronic feeding/oncogenicity study in rats with technical glyphosate. Document No. 008897. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. Available from: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/cleared-reviews/reviews/103601/103601-268.pdf>,

²⁷ Manas F, Peralta L, Raviolo J, Ovando HG, Weyers A, Ugnia L et al. (2009). Genotoxicity of glyphosate assessed by the comet assay and cytogenetic tests. *Environ Toxicol Pharmacol*, 28(1):37–41.

²⁸ Mladinic M, Berend S, Vrdoljak AL, Kopjar N, Radic B, Zeljezic D (2009). Evaluation of genome damage and its relation to oxidative stress induced by glyphosate in human lymphocytes in vitro. *Environ Mol Mutagen*, 50(9):800–7

²⁹ Alvarez-Moya C, Silva MR, Ramírez CV, Gallardo DG, Sánchez RL, Aguirre AC et al. (2014). Comparison of the in vivo and in vitro genotoxicity of glyphosate isopropylamine salt in three different organisms. *Genet Mol Biol*, 37(1):105–10.

³⁰ Monroy CM, Cortés AC, Sicard DM, de Restrepo HG (2005). Cytotoxicity and genotoxicity of human cells exposed in vitro to glyphosate. *Biomedica*, 25(3):335–45.

³¹ Koller VJ, Fürhacker M, Nersesyan A, Mišik M, Eisenbauer M, Knasmueller S (2012). Cytotoxic and DNA-damaging properties of glyphosate and Roundup in human-derived buccal epithelial cells. *Arch Toxicol*, 86(5):805–13.

³² Bolognesi C, Bonatti S, Degan P, Gallerani E, Peluso M, Rabboni R et al. (1997). Genotoxic activity of glyphosate and its technical formulation Roundup. *J Agric Food Chem*, 45(5):1957–62.

³³ Lioi MB, Scarfi MR, Santoro A, Barbieri R, Zeni O, Di Berardino D et al. (1998). Genotoxicity and oxidative stress induced by pesticide exposure in bovine lymphocyte cultures in vitro. *Mutat Res*, 403(1–2):13–20.

³⁴ Roustan A, Aye M, De Meo M, Di Giorgio C (2014). Genotoxicity of mixtures of glyphosate and atrazine and their environmental transformation products before and after photoactivation. *Chemosphere*, 108:93–100.

- 35 Moreno NC, Sofia SH, Martinez CB (2014). Genotoxic effects of the herbicide Roundup Transorb and its active ingredient glyphosate on the fish *Prochilodus lineatus*. *Environ Toxicol Pharmacol*, 37(1):448–54
- 36 Guilherme S, Gaivão I, Santos MA, Pacheco M (2010). European eel (*Anguilla anguilla*) genotoxic and pro-oxidant responses following short-term exposure to Roundup – a glyphosate-based herbicide. *Mutagenesis*, 25(5):523–30.
- 37 Lopes FM, Varela Junior AS, Corcini CD, da Silva AC, Guazzelli VG, Tavares G et al. (2014). Effect of glyphosate on the sperm quality of zebrafish *Daniorerio*. *Aquat Toxicol*, 155:322–6.
- 38 Paz-y-Mino C, Sanchez ME, Areval M, Munoz MJ, Witte T, De-la-Carrera GO et al. (2007). Evaluation of DNA damage in an Ecuadorian population exposed to glyphosate. *Genet Mol Biol*, 30(2):456–60.
- 39 Bolognesi C, Carrasquilla G, Volpi S, Solomon KR, Marshall EJ (2009). Biomonitoring of genotoxic risk in agricultural workers from five Colombian regions: association to occupational exposure to glyphosate. *J Toxicol Environ Health A*, 72(15–16):986–97.
- 40 Gehin A, Guillaume YC, Millet J, Guyon C, Nicod L (2005). Vitamins C and E reverse effect of herbicide-induced toxicity on human epidermal cells HaCaT: a biochemometric approach. *Int J Pharm*, 288(2):219–26.
- 41 Elie-Caille C, Heu C, Guyon C, Nicod L (2010). Morphological damages of a glyphosate-treated human keratinocyte cell line revealed by a micro- to nanoscale microscopic investigation. *Cell Biol Toxicol*, 26(4):331–9
- 42 Astiz M, de Alaniz MJ, Marra CA (2009b). Antioxidant defense system in rats simultaneously intoxicated with agrochemicals. *Environ Toxicol Pharmacol*, 28(3):465–73.
- 43 George J, Prasad S, Mahmood Z, Shukla Y (2010). Studies on glyphosate-induced carcinogenicity in mouse skin: a proteomic approach. *J Proteomics*, 73(5):951–64.
- 44 Thongprakaisang S, Thiantanawat A, Rangkadilok N, Suriyo T, Satayavivad J (2013). Glyphosate induces human breast cancer cells growth via estrogen receptors. *Food Chem Toxicol*, 59:129–36.
- 45 Gasnier C, Dumont C, Benachour N, Clair E, Chagnon MC, Séralini GE (2009). Glyphosate-based herbicides are toxic and endocrine disruptors in human cell lines. *Toxicology*, 262(3):184–91.
- 46 George J, Shukla Y (2013). Emptying of intracellular calcium pool and oxidative stress imbalance are associated with the glyphosate-induced proliferation in human skin keratinocytes HaCaT cells. *ISRN Dermatol*, 2013:825180

- 47 Chan P, Mahler J (1992). NTP technical report on the toxicity studies of glyphosate (CAS No. 1071-83-6) administered in dosed feed to F344/N rats and B6C3F1 mice. *Toxic Rep Ser*, 16:1-58.
- 48 Kreutz LC, Gil Barcellos LJ, de Faria Valle S, de Oliveira Silva T, Anziliero D, Davi dos Santos E et al. (2011). Altered hematological and immunological parameters in silver catfish (*Rhamdia quelen*) following short term exposure to sublethal concentration of glyphosate. *Fish Shellfish Immunol*, 30(1):51-7.
- 49 el-Gendy KS, Aly NM, el-Sebae AH (1998). Effects of edifenphos and glyphosate on the immune response and protein biosynthesis of boliti fish (*Tilapia nilotica*). *J Environ Sci Health B*, 33(2):135-49.
- 50 Reference Manual on Scientific Evidence, Third Edition, Committee on Science, Technology, and Law Policy and Global Affairs, FEDERAL JUDICIAL CENTER The National Academies Press.
- 51 Sperber, William H. (2001). "Hazard identification: from a quantitative to a qualitative approach". *Food Control*. 12: 223-228.
- 52 Ropeik, David (2002). *Risk*. New York, New York, USA: Houghton Mifflin Company.
- 53 Casarett & Doull 's *Toxicology: The Basic Science of Poisons* (8th edition). 2008. C.D. Klaassen (ed.). McGraw-Hill: New York. (Chapter 4).
- 54 NTP, RoC Process and Listing Criteria
<https://ntp.niehs.nih.gov/pubhealth/roc/process/index.html>
- 55 Huff JE, McConnell EE, Haseman JK, Boorman GA, Eustis SL, Schwetz BA, Rao GN, Jameson CW, Hart LG, Rall DP. Carcinogenesis studies: results of 398 experiments on 104 chemicals from the U.S. National Toxicology Program. *Ann NY Acad Sci*, 534:1-30.
- 56 Centers for Disease Control and Prevention. Principles of Epidemiology in Public Health Practice, Third Edition. An Introduction to Applied Epidemiology and Biostatistics. Available from:
<https://www.cdc.gov/ophss/csels/dsepd/ss1978/lesson1/section7.html>
- 57 Centers for Disease Control and Prevention. Principles of Epidemiology in Public Health Practice, Third Edition. An Introduction to Applied Epidemiology and Biostatistics. Available from:
<https://www.cdc.gov/ophss/csels/dsepd/ss1978/lesson1/section7.html>
- 58 Lee WJ, Cantor KP, Berzofsky JA, Zahm SH, Blair A (2004). Non-Hodgkin's lymphoma among asthmatics exposed to pesticides. *Int J Cancer*, 111(2):298-302

- ⁵⁹ Nordstrom M, Hardell L, Magnuson A, Hagberg H, Rask-Andersen A (1998). Occupational exposures, animal exposure and smoking as risk factors for hairy cell leukemia evaluated in a case-control study. *Br J Cancer*, 77(11):2048–52.
- ⁶⁰ Hardell L, Eriksson M (1999). A case-control study of non-Hodgkin lymphoma and exposure to pesticides. *Cancer*, 85(6):1353–60.
- ⁶¹ Orsi L, Delabre L, Monnereau A, Delval P, Berthou C, Fenaux P et al. (2009). Occupational exposure to pesticides and lymphoid neoplasms among men: results of a French case-control study. *Occup Environ Med*, 66(5):291–8.
- ⁶² Cocco P, Satta G, Dubois S, Pili C, Pilleri M, Zucca M et al. (2013). Lymphoma risk and occupational exposure to pesticides: results of the Epilymph study. *Occup Environ Med*, 70(2):91–8.
- ⁶¹ Karunanayake CP, Spinelli JJ, McLaughlin JR, Dosman JA, Pahwa P, McDuffie HH (2012). Hodgkin lymphoma and pesticides exposure in men: a Canadian case-control study. *J Agromed*, 17(1):30–9.
- ⁶² Brown LM, Blair A, Gibson R, Everett GD, Cantor KP, Schuman LM et al. (1990). Pesticide exposures and other agricultural risk factors for leukemia among men in Iowa and Minnesota. *Cancer Res*, 50(20):6585–91.
- ⁶³ Rank J, Jensen AG, Skov B, Pedersen LH, Jensen K (1993). Genotoxicity testing of the herbicide Roundup and its active ingredient glyphosate isopropylamine using the mouse bone marrow micronucleus test, Salmonella mutagenicity test, and phase-telophase test. *Mutat Res*, 300(1):29–36.
- ⁶⁴ <https://aghealth.nih.gov/about/index.html>
- ⁶⁵ Alavanja MC, Sandler DP, McMaster SB, Zahm SH, McDonnell CJ, Lynch CF et al. (1996). The Agricultural Health Study. *Environ Health Perspect*, 104(4):362–9
- ⁶⁶ De Roos AJ, Blair A, Rusiecki JA, Hoppin JA, Svec M, Dosemeci M et al. (2005). Cancer incidence among glyphosate-exposed pesticide applicators in the Agricultural Health Study. *Environ Health Perspect*, 113(1):49–54.
- ⁶⁷ De Roos AJ, Svec MA, Blair A, Rusiecki JA, Dosemeci M, Alavanja MC et al. (2005). Glyphosate results revisited. *Environ Health Perspect*, 113(6):A366–7
- ⁶⁸ Flower KB, Hoppin JA, Lynch CF, Blair A, Knott C, Shore DL et al. (2004). Cancer risk and parental pesticide application in children of Agricultural Health Study participants. *Environ Health Perspect*, 112(5):631–5.
- ⁶⁹ Engel LS, Hill DA, Hoppin JA, Lubin JH, Lynch CF, Pierce J et al. (2005). Pesticide use and breast cancer risk among farmers' wives in the Agricultural Health Study. *Am J Epidemiol*, 161(2): 121–35.

⁷⁰ Lee WJ, Sandler DP, Blair A, Samanic C, Cross AJ, Alavanja MC (2007). Pesticide use and colorectal cancer risk in the Agricultural Health Study. *Int J Cancer*, 121(2):339– 46.

⁷¹ Andreotti G, Freeman LE, Hou L, Coble J, Rusiecki J, Hoppin JA et al. (2009). Agricultural pesticide use and pancreatic cancer risk in the Agricultural Health Study Cohort. *Int J Cancer*, 124(10):2495–500

⁷² Dennis LK, Lynch CF, Sandler DP, Alavanja MC (2010). Pesticide use and cutaneous melanoma in pesticide applicators in the Agricultural Health Study. *Environ Health Perspect*, 118(6):812–7

⁷³ Schinasi L, Leon ME (2014). Non-Hodgkin lymphoma and occupational exposure to agricultural pesticide chemical groups and active ingredients: a systematic review and meta-analysis. *Int J Environ Res Public Health*, 11(4):4449–527.

⁷⁴ IARC (2015). International Agency for Research on Cancer. Monograph on Glyphosate. Volume 112

⁷⁵ Hill, Austin Bradford (1965). The Environment and Disease: Association or Causation?". *Proceedings of the Royal Society of Medicine*. 58 (5): 295–300.

⁷⁶ Knezevich, A.L and Hogan, G. K. (1983). A chronic feeding study of glyphosate in mice. Unpublished report prepared by Bio/Dynamic Inc., dated July 21, 1983. Report No. 77-2011. EPA Accession No. 251007 – 251009, and 251014

⁷⁷ EPA (2015) Glyphosate: Report of the Cancer Assessment Review Committee. EPA's Office of Pesticide Programs, Health Effects Division. October 1, 2015.

⁷⁸ Greim H, Saltmiras D, Mostert V, Strupp C (2015). Evaluation of carcinogenic potential of the herbicide glyphosate, drawing on tumor incidence data from fourteen chronic/carcinogenicity rodent studies. *Crit Rev Toxicol*, 45(3):185–208.

⁷⁹ EPA (1985b). EPA Reg.#: 524–308; Roundup; glyphosate; pathology report on additional kidney sections. Document No. 004855. Washington (DC): Office of Pesticides and Toxic Substances, United States Environmental Protection Agency. At: <http://www.epa.gov/pesticides/chemicalsearch/chemical/foia/clearedreviews/reviews/103601/103601-206.pdf>

⁸⁰ Chandra M, Frith CH (1994). Spontaneous renal lesions in CD-1 and B6C3F1 mice. *Exp Toxicol Pathol*, 46(3):189– 98.

⁸¹ EPA Memo, Glyphosate; EPA Registration No. 524-308; Roundup; Additional Histopathological Evaluations of Kidneys in the Chronic Feeding Study of Glyphosate in Mice, Toxicology Branch/HED. 11 Mar. 1986.

⁸² EPA Memo, Glyphosate EPA Reg. 524-308, 2 year chronic feeding/oncogenicity study in rats with technical glyphosate. 1986.

⁸³ Sauer, Robert M., *Pathology Working Group Report on Glyphosate in CD-1 Mice*, PATHCO, Inc., 10 Oct. 1985.

⁸⁴ Atkinson, C., Martin, T., Hudson, P., and Robb, D. (1993). Glyphosate: 104 week dietary carcinogenicity study in mice. Unpublished report prepared by Inveresk Research International April 7, 1993, Tranent, EH33 2NE, Scotland. IRI Project No. 438618. April 7, 1993. MRID 49631702.

⁸⁵ Sugimoto, K. (1997). 18-Month Oral Oncogenicity Study in Mice, Vol. 1 and 2. The Institute of Environmental Toxicology, 2-772, Suzuki-cho, Kodaira-shi, Tokyo, 187, Japan, Study No.: IET 94-0151.

⁸⁶ EPA (2016) Glyphosate Issue Paper: Evaluation of Carcinogenic Potential. EPA's Office of Pesticide Programs, September 12, 2016

⁸⁷ MONGLY02466412, Tier II Summaries for Glyphosate Carcinogenicity Studies from Greim et al., 2015 Paper, 9 November 2015.

⁸⁸ Kumar summary: Tables 20 and 21 from Feinchemie Schwebda 2001, Carcinogenicity Study with Glyphosate Technical in Swiss Albino Mice., unpublished, Study No.: Toxi: 1559. CARCI- M, Rallis India Ltd., Bangalore, India.

⁸⁹ David Saltmiras Deposition Ex. 5-29.

⁹⁰ Chruscielska K, Brzezinski J, Kita K, Kalhorn D, Kita I, Graffstein B et al. (2000). Glyphosate - Evaluation of chronic activity and possible far-reaching effects. Part 1. Studies on chronic toxicity. Pestycydy (Warsaw), 3-4:11-20.

⁹¹ Hard GC (1984). High frequency, single-dose model of renal adenoma/carcinoma induction using dimethylnitrosamine in Crl:(W) BR rats. Carcinogenesis 5:1047-1050

⁹² Lipsky ML and Trump BF (1988). Chemically induced renal epithelial neoplasia in experimental animals. Int. Rev. Exp. Pathol. 30:357-383

⁹³ Paz-y-Mino C, Sanchez ME, Areval M, Munoz MJ, Witte T, De-la-Carrera GO et al. (2007). Evaluation of DNA damage in an Ecuadorian population exposed to glyphosate. Genet Mol Biol, 30(2):456-60.

⁹⁴ Paz-y-Miño C, Muñoz MJ, Maldonado A, Valladares C, Cumbal N, Herrera C et al. (2011). Baseline determination in social, health, and genetic areas in communities affected by glyphosate aerial spraying on the northeastern Ecuadorian border. Rev Environ Health, 26(1):45-51.

- ⁹⁵ Sivikova K, Dianovsky J (2006). Cytogenetic effect of technical glyphosate on cultivated bovine peripheral lymphocytes Int J Hyg Environ Health, 209(1):15–20
- ⁹⁶ Guilherme S, Gaivão I, Santos MA, Pacheco M (2010). European eel (*Anguilla anguilla*) genotoxic and pro-oxidant responses following short-term exposure to Roundup—a glyphosate-based herbicide. Mutagenesis, 25(5):523–30.
- ⁹⁷ Gehin A, Guillaume YC, Millet J, Guyon C, Nicod L (2005). Vitamins C and E reverse effect of herbicide-induced toxicity on human epidermal cells HaCaT: a biochemometric approach. Int J Pharm, 288(2):219–26.
- ⁹⁸ Kwiatkowska M, Huras B, Bukowska B (2014). The effect of metabolites and impurities of glyphosate on human erythrocytes (in vitro). Pestic Biochem Physiol, 109:34–43
- ⁹⁹ Slaninova A, Smutna M, Modra H, Svobodova Z (2009). A review: oxidative stress in fish induced by pesticides. Neuro Endocrinol Lett, 30:Suppl 1: 2–12.
- ¹⁰⁰ Wang SS, Davis S, Cerhan JR, Hartge P, Severson RK, Cozen W, Lan Q, Welch R, Chanock SJ, and Rothman N. (2006) Polymorphisms in oxidative stress genes and risk for non-Hodgkin lymphoma. Carcinogenesis vol.27 no.9 pp.1828–1834, 2006
- ¹⁰¹ Lan Q, Zheng T, Shen M, Zhang Y, Wang SS, Zahm SH, Holford TR, Leaderer, Boyle P, Chanock S. (2007) Genetic polymorphisms in the oxidative stress pathway and susceptibility to non-Hodgkin lymphoma. Hum Genet 121:161–168

Studies I reviewed but determined inadequate for use:

Greim⁷⁸ reported on a study (Study 4, Suresh): Greim H, Saltmiras D, Mostert V, Strupp C (2015). Evaluation of carcinogenic potential of the herbicide glyphosate, drawing on tumor incidence data from fourteen chronic/carcinogenicity rodent studies. Crit Rev Toxicol, 45(3):185–208.

Greim⁷⁸ reported on a study (Study 8, Wood 2009a) : Greim H, Saltmiras D, Mostert V, Strupp C (2015). Evaluation of carcinogenic potential of the herbicide glyphosate, drawing on tumor incidence data from fourteen chronic/carcinogenicity rodent studies. Crit Rev Toxicol, 45(3):185–208.

Chruscielska K, Brzezinski J, Kita K, Kalhorn D, Kita I, Graffstein B et al. (2000). Glyphosate - Evaluation of chronic activity and possible far-reaching effects. Part 1. Studies on chronic toxicity. Pestycydy (Warsaw), 3–4:11–20.

Seralini GE, Clair E, Mesnage R, Gress S, Defarge N, Manuela Malatesta M et al. (2014). Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. Environmental Sciences Europe, 26(1):1–14

George J, Prasad S, Mahmood Z, Shukla Y (2010). Studies on glyphosate-induced carcinogenicity in mouse skin: a proteomic approach. J Proteomics, 73(5):951–64.

EXHIBIT A

C W Jameson - Curriculum Vitae and Bibliography

Name Charles William Jameson

Mailing Address: [REDACTED]
[REDACTED]

Date And Place Of Birth: [REDACTED]

Citizenship: [REDACTED]

Marital Status: Married, four children

Education: B.S. 1970
Chemistry,
Mount Saint Mary's College
Emmitsburg, Maryland

Ph.D. 1975
Organic Chemistry, Physical Chemistry minor
University of Maryland
College Park, Maryland

Brief Chronology of Employment:

1965	Chemistry Laboratory Technician, Bionetics Research Laboratories, Falls Church, Virginia
1968 – 1969:	Organic Chemistry Laboratory Assistant, Mount Saint Mary's College, Emmitsburg, Maryland
1969 – 1970:	Organic Chemistry Laboratory Instructor, Mount Saint Mary's College, Emmitsburg, Maryland
1970 – 1973:	Graduate Teaching Assistant, Chemistry Dept., University of Maryland College Park, Maryland
1973 – 1975:	Graduate Research Assistant, Center of Materials Research, University of Maryland, College Park, Maryland
1975 – 1976	Faculty Graduate Assistant, Chemistry Dept., University of Maryland, College Park, Maryland
1976 – 1979:	Senior Chemist, Tracor Jitco, Inc., Rockville, Maryland
1979 – 1980:	Chemist, Carcinogenesis Testing Program, National Cancer Institute, National Institutes of Health (NIH), Bethesda, Maryland

C W Jameson - Curriculum Vitae and Bibliography

1980 – 1983:	Head, Chemistry Section, Program Resources Branch, National Toxicology Program (NTP), National Institute of Environmental Health Sciences (NIEHS), NIH, Research Triangle Park, North Carolina
1983 – 1985:	Acting Chief, Program Resources Branch, NTP, NIEHS, NIH, Research Triangle Park, North Carolina
1985 – 1989:	Head, Program Resources Group, Carcinogenesis and Toxicologic Evaluation Branch, NTP, NIEHS, NIH, Research Triangle Park, North Carolina
1989 – 1990:	Supervisory Chemist, Experimental Toxicology Branch, NTP, NIEHS, NIH, Research Triangle Park, North Carolina
1990 – 1995:	Senior Chemist, Office of the Senior Scientific Advisor to the Director NIEHS, NIH, Research Triangle Park, North Carolina
1995 – 2008	Director, Report on Carcinogens, NTP, NIEHS, NIH, Research Triangle Park, North Carolina
2008 – present	Principal, CWJ Consulting, LLC, Cape Coral, Florida

Department of Health and Human Services Activities

Chairman, National Toxicology Program's Executive Committee's Interagency Working Group for the Report on Carcinogens, 1995 to 2005

National Institutes of Health Activities

NIEHS Representative to the Deafness and Other Communication Disorders Interagency Coordination Committee, 1990 - 1996.

NIEHS Representative on the Task Force on Aging Research, 1990-1994.

National Institutes of Environmental Health Sciences Activities

Chairman, NIEHS/NTP Review Committee for the Report on Carcinogens, 1995 to 2005

Chairman, Search Committee for NIEHS Tenure / Tenure Track Staff Epidemiologist 1998

Peer-Review Panel Member for Draft Report on Carcinogens Monograph on Cobalt and Certain Cobalt Compounds. July, 2015

Member and Chairman for the Special Emphasis Panel to review proposals responding to RFP ES2015038, "Scientific Information Management and Literature-Based Evaluations for the National

C W Jameson - Curriculum Vitae and Bibliography

Toxicology Program (NTP).” The objective of this contract is to provide scientific and technical expertise and support for the NTP to compile, review, and analyze information and data from the scientific literature and other sources regarding the effects of environmental substances and other issues that may impact public health. October, 2015

International Activities

Member, WHO Task Group on Environmental Health Criteria for Fully Halogenated Chlorofluorocarbons, Neuherberg, Federal Republic of Germany, November 21 – 25, 1988.

Member, WHO Task Group on Environmental Health Criteria for Partially Halogenated Chlorofluorocarbons (Ethane Derivatives), Carshalton, Surrey, United Kingdom, September 30 – October 5, 1991.

NIEHS representative to the WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 82 on the Carcinogenic Risks To Humans Of Some Traditional Herbal Medicines, Some Mycotoxins, Naphthalene And Styrene, Lyon, France, February 11 – 20, 2002

Member, IARC *Monographs* Advisory Group for Five Year Plan, Lyon, France, 18-21 February 2003

NIEHS representative to the WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 87 on The Carcinogenic Risks To Humans Of Lead And Lead Compounds, Lyon, France, February 8 – 18, 2004

NIEHS representative to the WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 91 on The Carcinogenic Risks To Humans Of Combined Oral Contraceptives And Estrogen-Progestogen Replacement Therapy, Lyon, France, June 4-15, 2005.

NIEHS representative to the WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 93 on The Carcinogenic Risks To Humans Of Carbon Black, Titanium Dioxide And Non-Asbestiform Talc, Lyon, France, February 4 – 15, 2006

Member, WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 97 on The Carcinogenic Risks To Humans Of 1,3 –Butadiene, Ethylene Oxide, And Vinyl Halides (Vinyl Fluoride, Vinyl Chloride And Vinyl Bromide), Lyon, France, June 6-15, 2007.

Member and Chair of Experimental Animal Data Subgroup, WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 99 on The Carcinogenic Risks To Humans Of Some Industrial And Cosmetic Dyes And Related Exposures, Lyon, France, February 4-13, 2008.

Member, WHO’s International Agency for Research on Cancer (IARC) Workgroup preparing Monograph 100A on A Review Of Human Carcinogens - Pharmaceuticals (Anti-Cancer Drugs – Hormonal Drugs & Therapies – Others), Lyon, France, October 14 – 21, 2008.

C W Jameson - Curriculum Vitae and Bibliography

Member and Chair of Experimental Animal Data Subgroup, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 100F on A Review Of Human Carcinogens - Chemical Agents And Related Occupations, Lyon, France, October 20 – 27, 2009.

Member and Chair of Experimental Animal Data Subgroup, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 103 on Bitumen And Bitumen Fumes, And Some Heterocyclic Aromatic Hydrocarbons, Lyon, France, October 11 - 18, 2011.

Member and Chair of Experimental Animal Data Subgroup, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 105 on Diesel And Gasoline Exhausts And Some Nitroarenes, Lyon, France, June 5 - 12, 2012.

Member WHO's International Agency for Research on Cancer (IARC) Workshop on Tumour Concordance And Mechanisms Of Carcinogenesis: Lessons Learned From Volume 100 of the IARC Monographs, Lyon, France: April 16-18, 2012 and November 28-30, 2012

Member and Chair of Experimental Animal Data Subgroup, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 108 On Some Drugs And Herbal Medicines, Lyon, France, June 4 - 11, 2013.

Member and Chair of Experimental Animal Data Subgroup, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 112 on Some Organophosphate Insecticides And Herbicides, Lyon, France, March 3-10, 2015.

Member and overall Chair, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph Vol. 115 on Some Industrial Chemicals, Lyon, France, February 2-9, 2016.

Member, WHO's International Agency for Research on Cancer (IARC) Workgroup preparing Monograph 116 on Coffee, Mate And Very Hot Beverages, Lyon, France, May 24 – 31, 2016.

Honors and Awards

President, Student Affiliate Chapter of the American Chemical Society, Mount Saint Mary's College, 1969; Vice President, 1968.

National Toxicology Program Representative to American Chemical Society's Committee on Regulatory Affairs 1982 – 1992.

National Institutes of Health Special Achievement Cash Award (Spy Dust Project): 1986.

Merit Pay Cash Award for Sustained High Quality Work Performance, NIEHS: 1982, 1989

Performance Award for Sustained High Quality Work Performance, NIEHS: 1991, 1992, 1993, 1995, 1996, 2001, 2002, 2003, 2004, 2006, 2007.

C W Jameson - Curriculum Vitae and Bibliography

Special Act or Service Award, NIEHS: 1996 (Review of Report on Carcinogens criteria); 1997 (Publication of 8th Report on Carcinogens); 1998 (Recruitment of NTP Staff Epidemiologist), 1998 (Restructuring of lead biokinetics contract and establishment of new Report on Carcinogens support contract)

Staff Recognition Award, NIEHS: 1999 (Preparation of final draft of 9th Report on Carcinogens)

NIEHS Director's Award, NIEHS: 2000 (Review of nominations for the 9th Report on Carcinogens)

Special Training

American Chemical Society, Short Course: "Chemical Carcinogenesis," 1978.

National Institutes of Health (NIH) Training Course: "Project Officers Civil Rights Contract Compliance," 1979.

Department of Health and Human Services Training (DHHS) Course: "Program Officials Guide to Contracting," 1980.

U. S. Office of Personnel Management (OPM) Training Course: "EEO - Its Place in the Federal Government," 1983.

U. S. OPM Training Course: "Introduction to Supervision," 1984.

NIH Training Course: "Employee Performance Management System Training," 1984.

DHHS Training Course: "Advanced Project Officer Training," 1985.

National Institute of Environmental Health Sciences Training Course: "Care and Handling of Laboratory Animals," 1986.

Rockhurst College Continuing Education Center: "How to Manage Projects, Priorities and Deadlines," 1992.

NIH Training Course: "PHS Animal Welfare Policy for HSA's," 1993.

Fred Pryor Seminars: "Total Quality Management," 1994.

Fred Pryor Seminars: "How to Manage Priorities and Meet Deadlines," 1994.

NIH Training Course: "Workplace Violence," 1994.

NIH Training Course: "NIH Guidelines on the Inclusion of Women and Minorities as Subjects in Clinical Research," 1994.

NIH Training Course: "Workplace Issues Associated with HIV/AIDS," 1994.

The Bookings Institution Course: "Issues in Science and Technology Policy", 1996

Professional Society Memberships and Activities

American Chemical Society

- Division of Analytical Chemistry
- Division of Chemical Health and Safety
- National Toxicology Program Representative to American Chemical Society's Committee on Regulatory Affairs 1982 – 1992
- Overall Co-Organizer and Co-Chairman of a symposium entitled "Chemistry and Safety for Toxicity Testing of Environmental Chemicals," sponsored by the Divisions of Chemical Health and Safety, Analytical Chemistry and Environmental Chemistry at the 183rd National American Chemical Society Meeting, Las Vegas, NV, March 1982.

Society of Toxicology

Research interests:

Chemical Carcinogenesis

Analytical chemistry methods development to support toxicology studies.

Reviewer for Scientific Journals

Analytical Chemistry

Bulletin of Environmental Contamination & Toxicology (Member of Editorial Board)

Environmental Health Perspectives (Contributing Editor)

Fundamental and Applied Toxicology

Journal of the National Cancer Institute

Science

Invited Papers

Invited to be Session Chairman and to present paper entitled "Analytical Chemistry Requirements for Toxicity Testing of Environmental Chemicals" at the Symposium on Chemistry and Safety for Toxicity Testing of Environmental Chemicals, at the 183rd National American Chemical Society Meeting, Las Vegas, NV, March 1982.

Invited to serve as a panelist on the NBC nationally televised series "Health Field" with Dr. Frank Field. A two-day series was filmed on Environmental Chemistry and Chemical Health Concerns, 1982.

C W Jameson - Curriculum Vitae and Bibliography

Invited to give a seminar entitled "Analytical Chemistry Requirements for Toxicity Testing." Duke University, Durham, NC, July 1982.

Invited to present a paper entitled "Practical Aspects of Analytical Chemistry Support for Toxicity Testing" at the Symposium on the Role of the Analytical Chemist in Animal and Molecular Toxicology, at the Federation of Analytical Chemistry and Spectroscopy Societies Meeting XI, Philadelphia, PA. September 16-21, 1984.

Invited to present a paper entitled "Application of Microencapsulation in Toxicity Testing" at the NIEHS Center Directors Meeting, Research Triangle Park, North Carolina, November 1984.

Invited to be Session Chairman and to present paper entitled "Chemical Quality Assurance Techniques for Toxicity Testing of Environmental Chemicals" at the Symposium on Accurate Measurements of Environmental Pollutants, at the 1984 International Chemical Congress of Pacific Basin Societies, Honolulu, Hawaii, December 16-21, 1984.

Invited to present a paper entitled "Lack of Evidence for Involvement of Cyanide in Methyl Isocyanate (MIC) Toxicity" at the Society of Toxicology Meeting, New Orleans, LA, March 3-7, 1986.

Invited to present a paper entitled "Toxicology From A Chemist's Viewpoint" at the Mount Saint Mary's College Science Alumni Homecoming, Emmitsburg, Maryland, October 23-26, 1986.

Invited to be Session Chairman and to present paper entitled "Application of Microencapsulation for Toxicity Studies" at the Symposium on Techniques for Microencapsulation of Chemicals at the 198th National Meeting of the American Chemical Society, Dallas, Texas, April 10-14, 1989.

Invited to be Session Chairman and to present paper entitled "Application of a Fischer Rat Leukemia Transplant Model as a Screen for the Leukemogenic Potential of Chemicals" at the International Symposium on Toxicology, Beijing, P. R. China, October 16-19, 1990.

Invited to present a paper entitled "Investigation of Alternative Vehicles for Use in Toxicology Research: Use of Microencapsulated and Molecular Encapsulated Chemicals in Toxicity Studies" at the Institute of Pharmacology and Toxicology, Academy of Military Medical Sciences, Beijing, P. R. China, October 20, 1990.

Invited to present a paper entitled "Toxicology and Carcinogenicity Studies of d- Limonene in Male and Female F344 Rats and B6C3F1 Mice" at the Symposium on Food Phytochemicals for Cancer Chemoprevention at the 204th National Meeting of the American Chemical Society, Washington, D.C., August 23-28, 1992.

Invited to be a Faculty Member and to present talk entitled " The National Toxicology Program's Report on Carcinogens " at the Toxicology Forum, Washington, DC, February 1995.

Invited to be a Faculty Member and to present talk entitled " The Report On Carcinogens (RoC): Status Of The Review Of The Criteria For Listing Substances In The RoC " at the Toxicology Forum, Washington, DC, February 1996.

C W Jameson - Curriculum Vitae and Bibliography

Invited to be a Faculty Member and to present talk entitled " Update of 1997 review of Nominations for the 9th Report on Carcinogens " at the Toxicology Forum, Washington, DC, February 1998.

Invited to be a Faculty Member and to present talk entitled " NTP Report on Carcinogens: History and the Process " at the Toxicology Forum, Aspen, CO, July 1999.

BIBLIOGRAPHY

Publications

1. Mazzocchi PH, Ammon HL, **Jameson CW**. Lanthanide Shift Reagents III: Errors Resulting from the Neglect of Angle Dependence, *Tetrahedron Letters*, 573, 1973.
2. **Jameson CW**. I. Study of Lanthanide shift Reagent - Substrate Interaction in Solution. II. Competitive Photochemical Type I and Type II Reactions of Amides and Imides. Dissertation Abstracts, 1975.
3. Ennis DM, Kramer A, Mazzocchi PH, **Jameson CW**, Bailey WJ. Synthetic N-Releasing Biodegradable Soil Conditioners I, *Hort Science*, 10, 505, 1975.
4. Ammon HL, Mazzocchi PH, Colicelli E, **Jameson CW**, Liu L. A Convenient Method for Mixing ²H and ¹³C Lanthanide Induced Shift (LIS) Calculations, A Technique for Facilitating ¹³C Assignments, *Tetrahedron Letters*, 1745, 1976.
5. Ennis DM, Kramer A, **Jameson CW**, Mazzocchi PH, Bailey WJ. Structural Factors Influencing the Biodegradation of Imides, *Appl Environ Microbiology*, 35, 51, 1978.
6. Murrill EA, Woodhouse EJ, Olin SS, **Jameson CW**. Carcinogenesis Testing and Analytical Chemistry, *Analytical Chemistry*, 52, 1188A, 1980.
7. Douglas JF, Hamm TE, **Jameson CW**, Mahar H, Stinson S, Whitmire CE. Monitoring Guidelines for the Conduct of Carcinogen Bioassays. US Department of Health and Human Services. DHHS Publication No. (NIH) 81-1774. Washington, DC, US Government Printing Office, 80 pp., 1981.
8. Dieter MP, Luster MI, Boorman GA, **Jameson CW**, Dean JH, Cox JW. Immunological and Biochemical Responses in Mice Treated with Mercuric Chloride, *Toxicol Appl Pharmacol*, 68, 218, 1983.
9. **Jameson CW**, Dunnick JK, Brown RD, Murrill EA. Chemical Characterization of Psoralens Used in the National Toxicology Program Research Projects, *National Cancer Institute Monograph*, 66, 103, 1984.
10. Timmons L, Cannon M, Grese D, Brown R, Haile C, Murrill E, **Jameson CW**. Identification of Chlorinated Phenyl and Phenoxy Substituted Dibenzodioxin, Dibenzofuran and Diphenyl Ether Homologs in Commercial Grade Pentachlorophenol, *Analytical Letters*, 17(A4), 277-296, 1984.

C W Jameson - Curriculum Vitae and Bibliography

11. Timmons L, Steel D, Cannon M, Grese D, Brown R, Murrill E, **Jameson CW**. Identification of Bromotertrachlorophenol in Commercial Pentachlorophenol Samples, *Journal of Chromatography*, V 314, 476-481, 1984.
12. Dunnick JK, **Jameson CW**, Benson JM. Toxicology and Carcinogenesis Studies of Nickel Oxide, Nickel Subsulfide and Nickel Sulfate. *Annals of Clinical and Laboratory Science*. V14.N5. 400-401, 1984.
13. Lamb JC, IV, **Jameson CW**, Choudury H, Gulati D K. Fertility Assessment by Continuous Breeding: Evaluation of Diethylstilbestrol and a Comparison of Results from Two Laboratories. *J Amer Coll Toxicol* 4, 173, 1985.
14. Thigpen JE, Liu LA, Richter CB, Lebetkin EH, Haseman JK, **Jameson CW**. The Comparative Estrogenic Activity of Semipurified, Certified, Standard and Open Formula Rodent Diets. *Laboratory Animal Science*, V35, N5, 526-527, 1985.
15. Kline DA, Hanna GR, Kuhn GO, Honaker CB, **Jameson CW**. Preparation and Stability of Animal Feed Mixtures Dosed with Rotenone, *J Asso Off Anal Chem*, Vol. 69, #4, 660-663, 1986.
16. **Jameson CW**, Moseman RF, Collins BJ, Hooper ND. Spy Dust: Methods for the Detection and Cleanup of a Chemical Tracking Agent. *Analytical Chemistry*, 58, 915A, 1986.
17. Agarwal DK, Eustis S, Lamb JC, **Jameson CW**, Kluwe WM. Influence of Dietary Zinc on Di(2-ethylhexyl)phthalate-Induced Testicular Atrophy and Zinc Depletion in Adult-Rats. *Toxicology and Applied Pharmacology*, V84, N1, 12-24, 1986.
18. Boorman GA, Hong HL, **Jameson CW**, Yoshitomi K, Maronpot, RP. Regression of Methyl Bromide Induced Forestomach Lesions in the Rat. *Toxicology and Applied Pharmacology*, 86, 131-139, 1986.
19. Collins B, Goehl TJ, **Jameson CW**, Kuhn G, Dux T. Analytical Methods for the Analysis of Microencapsulated Trichloroethylene in Corn Oil, Feed Dosage Formulations and Rat Whole Blood. *J. of Analytical Toxicology*, 10, 236, 1986.
20. **Jameson CW**, NTP Technical Report on the Toxicology and Carcinogenesis Studies of Tetrakis(hydroxymethyl)phosphonium sulfate (THPS) and Tetrakis(hydroxymethyl)phosphonium Chloride (THPC) in F344/N Rats and B6C3F1 Mice (Gavage Studies). NIH Publication No. 296, 1987.
21. Dunnick J K, **Jameson CW**, Montgomery CA. Subchronic Toxicity of Propantheline Bromide Administered in the Feed to Fischer 344/N Rats and B6C3F1 Mice. *Fundamental and Applied Toxicology*, V9, N3, 496-503, 1987.
22. Germolec DR, Burleson GR, **Jameson CW**, Ackermann MF, Lamm KR, Hayes HT, Luster MI. Depression of Natural-Killer Cell-Activity by Ochratoxin-A. *Environmental Health Perspectives*, V75, No. 5, 145-145, 1987.

C W Jameson - Curriculum Vitae and Bibliography

23. **Jameson CW**, Moseman RF, Hooper ND, Collins BJ. Spy Dust - Detecting a Chemical Tracking Agent. *Environmental Health Perspectives*, V75, No. 5, 143-143, 1987.
24. Melnick RL, **Jameson CW**, Goehl TJ. Application of Microencapsulation for Toxicology Studies - Stability, Bioavailability, and Toxicity of Microencapsulated Trichloroethylene. *Environmental Health Perspectives*, V75, No. 5, 142-142, 1987.
25. Melnick RL, **Jameson CW**, Goehl TJ, Kuhn GO. Application of Microencapsulation for Toxicology Studies. 1. Principles and Stabilization of Trichloroethylene In Gelatin-Sorbitol Microcapsules. *Fundamental and Applied Toxicology*, V8, N4, 425-431, 1987.
26. Melnick RL, **Jameson CW**, Goehl TJ, Maronpot RR, Collins BJ, Greenwell A, Harrington FW, Wilson RE, Tomaszewski KE, Agarwal DW. Application of Microencapsulation for Toxicology Studies. 2. Toxicity of Microencapsulated Trichloroethylene in Fischer 344 Rats. *Fundamental and Applied Toxicology*, V8, N4, 432-442, 1987.
27. Thigpen JE, Lung-An L, Richter CB, Lebetkin, EH, Haseman, JK, **Jameson CW**. The Mouse Bioassay Test for the Detection of Estrogenic Activity in Feeds and Foodstuffs. Part I: A Standardized Method for Conducting the Mouse Bioassay using the CD-1 Mouse. *Laboratory Animal Science*, V37, N5, 596-601, 1987.
28. Thigpen JE, Lung-An L, Richter CB, Lebetkin EH, **Jameson CW**. The Mouse Bioassay Test for the Detection of Estrogenic Activity in Feeds and Foodstuffs. Part II: The Comparative Estrogenic Activity of Purified, Certified Standard, Open and Closed Formula Rodent Diets. *Laboratory Animal Science*, V37, N5, 602-605, 1987.
29. Bucher JR, Gupta BN, Adkins B, Thompson M, **Jameson CW**, Thigpen J E, Schwetz BA. The Toxicity of Inhaled Methyl Isocyanate in F344/N Rats and B6C3F1 Mice. I: Acute Exposure and Recovery Studies. *Environmental Health Perspectives*, V72, 53-61, 1987.
30. Luster MI, Germolec DR, Burleson GR, **Jameson CW**, Ackermann MF, Lamm KR, Hayes HT. Selective Immunosuppression in Mice of Natural Killer Cell Activity by Ochratoxin A. *Cancer Research*, Vol. 47, 2259-2263, 1987.
31. Dieter MP, **Jameson CW**, Tucker AN, Luster MI, French JE, Hong, HL, Boorman, GA. Evaluation of Tissue Disposition, Myelopoietic and Immunologic Responses in Mice After Long-term Exposure to Nickel Sulfate in the Drinking Water. *Journal of Toxicology and Environmental Health*, V24, 357-372, 1988.
32. Huff JE, McConnell EE, Haseman JK, Boorman GA, Eustis SL, Schwetz BA, Rao GN, **Jameson CW**, Hart LG, Rall DP. Carcinogenesis Studies Results of 398 Experiments on 104 Chemicals from the U. S. National Toxicology Program. *Annals of the New York Academy of Sciences* V534, 1-30, 1988.
33. Shan A, Harben D, **Jameson CW**. Analyses of Two Azo Dyes by High Performance Liquid Chromatography. *Journal of Chromatographic Science*, V26, 439-442, 1988.

C W Jameson - Curriculum Vitae and Bibliography

34. Hong HL, Canipe J, **Jameson CW**, Boorman GA: Comparative Effects of Ethylene Glycol and Ethylene Glycol Monomethyl Ether Exposure on Hematopoiesis and Histopathology in B6C3F1 Mice. *Journal of Environmental Pathology, Toxicology, and Oncology*, V8, N7, 27-38, 1988.
35. Hong HL, **Jameson CW**, Boorman GA. Residual Hematopoietic Effect of Ochratoxin A in Mice Exposed to Irradiation. *Toxicology*, V53, 57-67, 1988.
36. Dieter MP, **Jameson CW**, French JE, Gangjee S, Stefanski SA, Chan, PC. Development and Validation of a Cellular Transplant Model for Leukemia in Fischer Rats: A Short-term Assay for Potential Anti-leukemic Chemicals. *Leukemia Research*, V13, 841-849, 1989.
37. Timmons L, Brown R, Arneson DW, **Jameson CW**. Rapid Determination of Low pg/mg Amounts of N-Nitrosodiethylamine in Rodent Body Fluid and Tissue Samples by Isotope-Dilution High Resolution Mass Spectrometry. *J. Anal. Tox.*, V13, N6, 333-336, 1989.
38. Heindel JJ, Lamb JC, Chapin RE, Gulati DK, Hope E, George J, **Jameson CW**, Teague J, Schwetz BA. Reproductive Toxicity Testing by Continuous Breeding Test Protocol in CD-1 Mice. DHHS Publication No. (NIH) 89 Washington, DC, US Government Printing Office, 1989.
39. Cannon JM, Brown D, Murrill EM, **Jameson CW**. Identification of Components in Iodinated Glycerol. *Journal of Pharmaceutical Sciences*, V78, N1, 48-51, 1989.
40. Morgan DL, **Jameson CW**, Mennear JH, Prejean JD. 14-Day and 90-Day Toxicity Studies of C.I. Pigment Red 3 in Fischer 344 Rats and B6C3F1 Mice. *Fd. Chem. Toxic.*, V27, N12, 793-800, 1989.
41. Morgan DL, **Jameson CW**, Mennear JH, Ulland BM. Thirteen-Week Toxicity Studies of CI Direct Blue 15 and 3,3'-Dimethoxybenzidine in the Fischer 344 Rat. *Toxicology*, V59, 297-309, 1989.
42. Dieter MP, **Jameson CW**, Maronpot RR, Langenbach RJ, Braun AG. The Chemotherapeutic Potential of Glycol Alkyl Ethers: Structure-Activity Studies of Nine Compounds in a Fischer Rat Leukemia Transplant Model. *Cancer Chemother. Pharmacol.*, 26, 173-180, 1990.
43. Gorski T, Goehl TJ, **Jameson CW**, Collins BJ. Sources of Error in the Determination of Trichloroethylene in Blood. *Bull. Environ. Contam. Toxicol.*, V45, 1-5, 1990.
44. Dieter MP, Boorman GA, **Jameson CW**, Matthews HB, Huff JE. The Carcinogenic Activity of Commercial Grade Toluene Diisocyanate in Rats and Mice in Relation to the Metabolism of the 2,4- and 2,6-TDI Isomers. *Toxicology and Industrial Health*, V6, No. 6, 599-621, 1990.
45. Morrissey RE, Fowler BA, Harris MA, Moorman MP, **Jameson CW**, Schwetz BA. Arsine: Absence of Developmental Toxicity in Rats and Mice. *Fundamental and Applied Toxicology* 15, 350-356, 1990.
46. **Jameson CW**, NTP Technical Report on the Toxicology and Carcinogenesis Studies of d-Limonene in F344/N Rats and B6C3F1 Mice (Gavage Studies). NIH Publication No. 347, 1990.
47. Gorski T, Goehl TJ, **Jameson CW**, Collins BJ, Bursey J, Moseman R. Gas Chromatic Determination of 2-Ethylhexanol and 2-Ethylhexanoic Acid as Derivatives suitable for Electron Capture and Nitrogen-

C W Jameson - Curriculum Vitae and Bibliography

Phosphorus Detection After Single Reaction with Heptafluorobutyrimidazole. *Journal of Chromatography*, 509, 383-389, 1990

48. Yuan J, **Jameson CW**, Goehl TJ, Collins BJ, Corniffee G, Kuhn G, Castro C. Effects of Physical Binding of o-Nitroanisole with Feed Upon its Systemic Availability in Male F344 Rats. *Bulletin of Environmental Contamination and Toxicology*, 47: 152-159, 1991.
49. Yuan J, **Jameson CW**, Goehl TJ, Collins BJ, Purde W, Judd L. Application of Molecular Encapsulation for Toxicity Studies: Toxicokinetics of p-Chloro- α,α,α -trifluorotoluene in β -Cyclodextrin or Corn Oil Vehicles in Male F344 Rats. *Toxicology and Applied Pharmacology*, 111, 107-115, 1991.
50. Dieter MP, **Jameson CW**, Elwell M, Lodge JW, Hejtmancik M, Grumbein SL, Ryan M, Peters AC. Comparative Toxicity and Tissue Distribution of Antimony Potassium Tartrate in Rats and Mice Dosed by Drinking Water and Intraperitoneal Injection. *Journal of Toxicology and Environmental Health*, 34, 51-82, 1991.
51. Yuan J, Bucher JR, Goehl TJ, Dieter MP, **Jameson CW**. Quantitation of Cinnamaldehyde and Cinnamic Acid in Blood by HPLC. *Journal of Analytical Toxicology*, 16, N6: 359-362, 1992.
52. Yuan J, **Jameson CW**, Goehl TJ, Elwell MR, Leininger JR, Thompson MB, Corniffe G, Carleton T. Application of Molecular Encapsulation for Toxicology Studies: Comparative Toxicity of p-Chloro- α,α,α -trifluorotoluene in β -Cyclodextrin Vehicle versus Corn Oil Vehicle in Male and Female Fischer 344 Rats and B6C3F1 Mice. *Fundamental and Applied Toxicology*, 18, 460-470, 1992.
53. Dieter MP, Maronpot RR, **Jameson CW**, Ward SM. The Effects of Iodinated Glycerol, Trichlorfon, Acetaminophen on Tumor Progression in a Fischer Rat Leukemia Transplant Model. *Cancer Detection and Prevention*, V16, No. 3, 173-183, 1992.
54. Yuan J, Dieter MP, Bucher JR, **Jameson CW**. Toxicokinetics of Cinnamaldehyde in F344 Rats. *Food and Chemical Toxicology*, 30, N12: 997-1004, 1992.
55. Yuan J, **Jameson CW**, Goehl TJ, Collins BJ. Molecular Encapsulator: A Novel Vehicle for Toxicology Studies. *Toxicology Methods*, V1, No.4, 231-241, 1992.
56. Dieter MP, Boorman GA, **Jameson CW**, Eustis SL. Development of Renal Toxicity in F344 Rats gavaged with Mercuric Chloride for 2 Weeks, or 2, 4, 6, 15, and 24 Months. *Journal of Toxicology and Environmental Health*, 36, 319-340, 1992.
57. Yuan J, Dieter MP, Bucher JR, **Jameson CW**. Application of Microencapsulation for Toxicology Studies III, Bioavailability of Microencapsulated Cinnamaldehyde. *Fundamental and Applied Toxicology*, 20, N1: 83-87, 1993.
58. Dieter MP, Goehl TJ, **Jameson CW**, Elwell MR, Hildebrandt PK, Yuan J. Comparison of the Toxicity of Citral in F344 Rats and B6C3F1 Mice When Administered by Microencapsulation in Feed or by Corn Oil Gavage. *Food and Chemical Toxicology*, 31, N7: 463-474, 1993.

C W Jameson - Curriculum Vitae and Bibliography

59. Arneson DA, Kuhn GO, **Jameson CW**. Analysis of Feed Blends Containing Microencapsulated 2-Ethyl-1-hexanol: Verification of Homogeneity and Stability. *Journal of Applied Toxicology*, 15 (1), 1-4, 1995.
60. **Jameson CW**, Ed. Conference on Beryllium Related Diseases. *Environmental Health Perspectives*, Vol. 104, S5, 935-998, 1996.
61. **Jameson CW**. Introduction to the Conference on Beryllium Related Diseases. *Environmental Health Perspectives*, Vol. 104, S5, 935-936, 1996.
62. Gulson BL, **Jameson CW**, Mahaffey KR, Mizon KJ, Korsch MJ, Vimpani, G. Pregnancy increases mobilization of lead from maternal skeleton. *J Lab Clin Med.*, 130, 51-62, 1997.
63. Gulson BL, Gillings BR, **Jameson CW**. Stable lead isotopes in teeth as indicators of past domicile - a potential new tool in forensic science. *J Forensic Sciences*, 42, 787-791, 1997.
64. Gulson BL, Mahaffey KR, **Jameson CW**, Vidal M, Law AJ, Mizon KJ, Korsch MJ. Dietary Intake for Mother-Child Pairs and Implications for Pharmacokinetic Models. *Environ Health Persp.*, 105, 1334-1342, 1997.
65. Gulson BL, Cameron MA, Smith AJ, Mizon KJ, Korsch MJ, Vimpani G, McMichael AJ, Pisaniello D, **Jameson CW**, Mahaffey KR. Blood lead-urine relationships in adults and children. *Environ Res. Section A* 78, 152-160, 1998.
66. Gulson BL, **Jameson CW**, Mahaffey KR, Mizon KJ, Korsch MJ, Cameron MA, Eisman JA. Mobilization of lead from the skeleton during the post-natal period is larger than during pregnancy. *J Lab Clin Med.*, 131, 324-329, 1998.
67. Gulson BL, **Jameson CW**, Mahaffey KR, Mizon KJ, Patison N, Law JL, Korsch MJ, Salter MA. Relationship of Lead in Breast Milk to Lead in Blood, Urine, and Diet of the Infant and Mother. *Environ Health Persp.*, 106, 667-674, 1998.
68. Gulson BL, Gray B, Mahaffey KR, **Jameson CW**, Mizon KJ, Patison N, Korsch MJ. Comparison of the rates of exchange of lead in the blood of newly born infants and their mothers with lead from their current environment, *J Lab Clin Med.*, 133, Vol. 2, 171-178, 1999.
69. Gulson BL, Mahaffey KR, **Jameson CW**, Patison N, Law JL, Mizon KJ, Korsch MJ, Pederson, D. Impact of Diet on Lead in Blood and Urine in Female Adults and Relevance to Mobilization of Lead from Bone Stores. *Environ Health Persp.*, 107, N4, 257-263, 1999.
70. Bucher JR, **Jameson CW**. Environmental tobacco smoke epidemiology. *Environ Health Perspect.* 107(8): A395, 1999.
71. Waalkes, M. P. and **Jameson, C. W.**: Evaluation of nickel compounds for listing in the Report on Carcinogens. In: Vernet, P. G. (ed.). *Proceedings of the Sixth International Symposium on Metal Ions in Biology and Medicine*. Montrouge, France, John Libby Eurotext, Ltd. In press, 2000.

C W Jameson - Curriculum Vitae and Bibliography

72. Gulson BL, Mizon KJ, Palmer JM, Korsch MJ, Patison N, **Jameson CW**, Donnelly JB. Urinary lead isotopes during pregnancy and postpartum indicate no preferential partitioning of endogenous lead into plasma. *J Lab Clin Med.*, 136(3): 236-42, 2000.
73. Portier CJ, et al. Differences in the carcinogenic evaluation of glyphosate between the International Agency for Research on Cancer (IARC) and the European Food Safety Authority (EFSA) *J Epidemiol Community Health* August Vol 70 No 8: 741-745, 2016.

Listed as a contributor for the evaluation, interpretation and reporting of results for more than 100 chemicals studied in chronic two-year bioassay studies by the National Toxicology Program as published in the Technical Report Series (1980-1990).

BOOKS

Jameson CW and Walters DB, Eds. *Chemistry for Toxicity Testing*, Butterworth Publishers, Boston, MA, 1984.

Walters DB and **Jameson CW**, Eds. *Health and Safety for Toxicity Testing*, Butterworth Publishers, Boston, MA, 1984.

REPORTS

Jameson CW, Editor, *Report on Carcinogens*, Eighth Edition, U.S. Department of Health and Human Services, Public Health Service, 1998.

Jameson CW, Editor, *Report on Carcinogens*, Ninth Edition, U.S. Department of Health and Human Services, Public Health Service, 2000.

Jameson CW, Editor, *Report on Carcinogens*, Tenth Edition, U.S. Department of Health and Human Services, Public Health Service, 2002.

Jameson CW, Editor, *Report on Carcinogens*, Eleventh Edition, U.S. Department of Health and Human Services, Public Health Service, 2004.

Jameson CW, Senior Author for following NTP Report on Carcinogens Background Documents:

1. Alcoholic Beverage Consumption - 1999
2. 1-Amino-2,4-dibromoanthraquinone - 2002
3. 2-Amino-3,4-dimethylimidazo[4-5-f]quinoline (MeIQ) - 2002
4. 2-Amino-3,8-dimethylimidazo[4-5-f]quinoxaline (MeIQx) - 2002
5. 2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) - 2002
6. 2-Amino-3-methylimidazo[4,5-f]quinoline (IQ) - 2002
7. Azacitidine - 1996
8. Beryllium and Beryllium Compounds - 2000

C W Jameson - Curriculum Vitae and Bibliography

9. 2,2-bis-(bromomethyl)-1,3-propanediol (BBMP) (Technical Grade) - 2000
10. Boot & Shoe Manufacturing - 1998
11. 1,3- Butadiene - 1997
12. Cadmium and Cadmium Compounds - 1997
13. Chloramphenicol - 2000
14. Chloroprene - 1997
15. Chlorozotocin - 1996
16. p -Chloro-o-toluidine and its Hydrochloride Salt - 1996
17. Cobalt Sulfate - 2002
18. Cyclosporin A - 1996
19. Danthron (1,8-Dihydroxyanthraquinone) - 1996
20. Diazoaminobenzene - 2002
21. 2,3-Dibromo-1-propanol - 2000
22. Diesel Exhaust Particulates - 1998
23. Diethanolamine - 2002
24. 1,6-Dinitropyrene & 1,8-Dinitropyrene - 1996
25. Disperse Blue I - 1996
26. Dyes Metabolized to Benzidine (Benzidine Dyes as a Class) - 1997
27. Dyes metabolized to 3,3'-Dimethoxybenzidine (DMOB) - 2000
28. Dyes metabolized to 3,3'-Dimethylbenzidine (DMB) - 2000
29. Environmental Tobacco Smoke - 1998
30. Estrogens, Steroidal - 2000
31. Ethyl Acrylate - 1998
32. Ethylene Oxide - 1998
33. Furan - 1996
34. Hepatitis B Virus (HBV) - 2003
35. Hepatitis C Virus (HCV) - 2003
36. Human Papillomaviruses (HPV): Some Genital-Mucosal Types - 2003
37. Isoprene - 1998
38. Lead and Lead Compound - 2003
39. Methyleugenol - 2000
40. Methyl-t-Butyl Ether (MtBE) - 1998
41. Naphthalene - 2002
42. Nickel Compounds - 1998
43. Nickel (Metallic) and Certain Nickel Alloys - 2000
44. o-Nitroanisole - 1996
45. Nitrobenzene - 2002
46. 6-Nitrochrysene - 1996
47. Nitromethane - 2002
48. 1-Nitropyrene - 1996
49. 4-Nitropyrene - 1996
50. Phenolphthalein - 1997
51. Saccharin - 1997
52. Silica, Crystalline (Respirable Size) - 1998
53. Smokeless Tobacco - 1997
54. Solar Radiation & Exposure to Sunlamps or Sunbeds - 1997
55. Strong Inorganic Acid Mists Containing Sulfuric Acid - 1997
56. Styrene-7,8-oxide - 2000

C W Jameson - Curriculum Vitae and Bibliography

57. Tamoxifen - 1997
58. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) - 1997
59. Tetrafluoroethylene - 1997
60. 4,4'-Thiodianiline - 2002
61. Thiotepa - 1996
62. Tobacco Smoking - 1997
63. 1,2,3-Trichloropropane - 1996
64. Trichloroethylene - 1997, 2000
65. Ultraviolet (UV) Radiation, Broad Spectrum and UVA, UVB, and UVC - 2000
66. Vinyl Bromide - 2000
67. Vinyl Fluoride - 2000
68. Wood Dust - 2000
69. X-Radiation & Gamma Radiation and Neutrons – 2003

Jameson CW Contributor to the following NTP Report on Carcinogens Background Documents:

1. Aristolochic Acid Related Exposures (2 Candidate Substances) - 2008
 - a. Botanical Products Containing Aristolochic Acid
 - b. Aristolochic Acid
2. Captafol – 2008
3. ortho-Nitrotoluene – 2008
4. Riddelliine - 2008
5. Styrene - 2008
6. Cobalt – Tungsten Carbide; Powders and Hard Metals – 2009
7. Glass Wool Fibers – 2009

BOOK CHAPTERS

1. **Jameson CW**. Analytical Chemistry Requirements for Toxicity Testing of Environmental Chemicals, in Chemistry for Toxicity Testing, pp. 3-14, Butterworth Publishers, Boston, MA, 1984.
2. **Jameson CW**, Rollheiser JJ, Kuhn GO. Stability Determinations of Chemical/Vehicle Mixtures, in Chemistry for Toxicity Testing, pp. 107-114, Butterworth Publishers, Boston, MA, 1984.
3. Kuhn GO, Rollheiser JJ, Schworer BA, **Jameson CW**. Methods Development for Mixing Chemicals in Rodent Feed, in Chemistry for Toxicity Testing, pp. 59-81, Butterworth Publishers, Boston, MA, 1984.
4. Murrill EA, Kuhn GO, Rollheiser JJ, **Jameson CW**. Analysis of Dose Feed Mixtures, in Chemistry for Toxicity Testing, pp. 91-106, Butterworth Publishers, Boston, MA, 1984.
5. Woodhouse EJ, Murrill EA, Stelting KM, **Jameson CW**. Problems of Testing Commercial-Grade Chemicals, in Chemistry for Toxicity Testing, pp. 31-50, Butterworth Publishers, Boston, MA, 1984.

C W Jameson - Curriculum Vitae and Bibliography

6. Graves SW, Woodhouse EJ, Stelting KM, **Jameson CW**: Bulk Chemical Management for chronic Toxicity Studies, in Health and Safety for Toxicity Testing, pp. 221-240, Butterworth Publishers, Boston, MA, 1984.
7. Huff JE, McConnell EE, Haseman JK, Boorman GA, Eustis SL, Schwetz BA, Rao GA, **Jameson CW**, Hart LG, Rall DP. Carcinogenesis Studies: Results of 398 Experiments on 104 Chemicals from the U. S. National Toxicology Program, in Living in a Chemical World, Occupational and Environmental Significance of Industrial Carcinogens, Annals of the New York Academy of Sciences, V 534, pp. 1-31, 1988.
8. Fouts JR, **Jameson CW**. Hazard Identification, The First Step, in Proceedings of the National Minority Health Conference, Atlanta GA, December 1990.
9. **Jameson CW**, Goehl TJ. Chemistry Requirements for the Toxicologic and Carcinogenicity Evaluation of Chemicals, in the Handbook of Carcinogen Testing, Second Edition, pp. 286-297, Noyes Publications, Park Ridge, NJ, 1994.

ABSTRACTS/PRESENTATIONS

1. Mazzocchi PH, **Jameson CW**, Nishiyama T. Competing Processes in the Photochemistry of Alkyl Imides. Proceedings of the 178th National Meeting of the American Chemical Society, Washington, DC, September 1979.
2. **Jameson CW**. An Overview of Analytical Chemistry Requirements for Toxicity Testing. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
3. **Jameson CW**, Rollheiser JJ, Kuhn GO. Stability Determinations of Chemical/Vehicle Mixtures. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
4. **Jameson CW**, Grieshaber CK, Whitmire CE. Effect of GLPs on Chemistry Requirements for Toxicity Testing. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
5. Kuhn GO, Rollheiser JJ, Schworer BA, **Jameson CW**. Methods Development for Mixing Chemicals in Rodent Feed. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
6. Murrill EA, Kuhn GO, Rollheiser JJ, **Jameson CW**. Analysis of Dose Feed Mixtures. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
7. Woodhouse EJ, Murrill EA, Brown RD, **Jameson CW**. The Problems of Testing Commercial Grade Chemicals. Proceedings of the 183rd National Meeting of the American Chemical Society, NV, March 1982.

C W Jameson - Curriculum Vitae and Bibliography

8. Graves SW, Woodhouse EJ, Stelting KM, **Jameson CW**. Bulk Chemical Management for Chronic Toxicity Studies. Proceedings of the 183rd National Meeting of the American Chemical Society, Las Vegas, NV, March 1982.
9. **Jameson CW**, Castro CA, Kuhn GO, Murrill EA. Quality Assurance Techniques for Reference laboratory Analysis of Dosage Formulations. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
10. Pallas FE, DuSold DE, Murrill EA, **Jameson CW**. HPLC Determination of Benzidine and its Congeners in Direct Dyes. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
11. Rollheiser JJ, Stelting KM, Woodhouse EJ, **Jameson CW**. Microcomputer Applications in a Bioanalytical Chemistry Laboratory. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
12. Fanska CB, Pittman LW, Murrill EA, **Jameson CW**, Dieter MP. HPLC Determination of Benzo(a)pyrene and Benzo(e)pyrene in Mouse Tissues. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
13. Cannon JM, Wyatt LL, Brown RD, Murrill EA, **Jameson CW**, Dieter MP. Analysis of Titanium in Tissues by IP Spectroscopy. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
14. Pittman LW, Lillich MA, Walters KM, Murrill EA, **Jameson CW**. HPLC Purity Analysis of Benzidine Congener-Based Dyes. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
15. Brown RD, Murrill EA, Stelting KM, Woodhouse EJ, **Jameson CW**. Selection of Analytical Methods for Characterization of Commercial Chemicals. Proceedings of the 184th National Meeting of the American Chemical Society, Kansas City, MO, September 1982.
16. Dieter MP, Luster MI, Dean JH, Boorman GA, **Jameson CW**. Immunotoxicity of Mercuric Chloride in B6C3F1 Mice. *The Toxicologist*, 2, 92, 1982.
17. Rollheiser JJ, Stelting KM, Woodhouse EJ, Kuhn GO, **Jameson CW**. A Systematic Procedure for the Development of Chromatographic Methods of Analysis of Dosed Feed Mixtures. Proceedings of the 184th National Meeting of the American Chemical Society, Seattle, WA, March 1983.
18. Minor CM, Graves SW, Stelting KM, Woodhouse EJ, **Jameson CW**. Microcomputer Uses in Quality Assurance. Proceedings of the 184th National Meeting of the American Chemical Society, Seattle, WA, March 1983.
19. Fanska CB, Shan A, Murrill EA, **Jameson CW**. HPLC Determination of Chloramphenicol and its Monosuccinate Esters in Rat and Mouse Sera. Proceedings of the 186th National Meeting of the American Chemical Society, Washington, DC, August 1983.

C W Jameson - Curriculum Vitae and Bibliography

20. Kuhn GO, Stelting KM, Dux T, **Jameson CW**. Analysis of o-Nitroanisole in Aged Feed Blends. Proceedings of the 186th National Meeting of the American Chemical Society, Washington, DC, August 1983.
21. Kuhn GO, Stelting KM, Kline DA, Murrill EA, **Jameson CW**. Determination of Sub-PPB Levels of Diethylstilbestrol in Feeds. Proceedings of the 184th National Meeting of the American Chemical Society, Washington, DC, August 1983.
22. Cannon JM, Brown RD, Fanska C, Woodhouse EJ, Murrill EA, **Jameson CW**. Chemical Characterization of a Commercial Dye for Use in Bioassay Studies. Proceedings of the 186th National Meeting of the American Chemical Society, Washington, DC, August 1983.
23. Timmons L, Cannon M, Grese D, Brown RD, Murrill EA, **Jameson CW**. Identification of Chlorinated Phenyl Ethers of Dibenzodioxins and Diphenyl Ethers in Commercial Grade Pentachlorophenol. Proceedings of the 186th National Meeting of the American Chemical Society, Washington, DC, August 1983.
24. Melnick RL, **Jameson CW**, Goehl TJ, Kuhn GO. Microencapsulation of Chemicals for Toxicologic Studies. *The Toxicologist*, 4, 49, 1984.
25. Onstot J, Timmons L, Murrill E A, **Jameson CW**. HRGC/HRMS Identification of Chlorinated Phenyl and Phenoxy Substituted Dioxins, Furans and Diphenyl Ethers in Commercial Grade Pentachlorophenol. Proceedings of the American Society for Mass Spectrometry Symposium, San Antonio, TX, May 1984.
26. Rollheiser J, Buchanan RC, Maune C, **Jameson CW**. Determination of Salicylazosulfapyridine in Rodent Feed. Proceedings of the 189th National Meeting of the American Chemical Society, Philadelphia, PA, August 1984.
27. Kuhn GO, Ridlen RL, Dix TP, Stelting, KM, **Jameson CW**. Conformance Testing of Microencapsulated Chemicals for Use in Bioassay Studies. Proceedings of the 189th National Meeting of the American Chemical Society, Philadelphia, PA, August 1984.
28. Kuhn GO, Arneson D, Kline DA, Shan YA, Nguyen PTH, **Jameson CW**. Determination of Metabolites of Benzidine-Based Dyes in Rodent Urine. Proceedings of the 189th National Meeting of the American Chemical Society, Philadelphia, PA, August 1984.
29. Dunnick JK, **Jameson CW**, Benson JM. Toxicology and Carcinogenesis Studies of Nickel Oxide, Nickel Subsulfide and Nickel Sulfate. Proceedings of the Third International Conference on Nickel Metabolism and Toxicology, Paris, France, September 1984.
30. **Jameson CW**. Practical Aspects of Analytical Chemistry Support for Toxicity Studies. Proceedings of the XI Federation of Analytical Chemistry and Spectroscopy Societies, Meeting, Philadelphia, PA, September 1984.

C W Jameson - Curriculum Vitae and Bibliography

31. Brown RD, Arneson DW, Stelting KM, **Jameson CW**. The Importance of Complementary Techniques in the Analysis of Commercial Chemicals for Toxicity Testing. Proceedings of the XI Federation of Analytical Chemistry and Spectroscopy Societies Meeting, Philadelphia, PA, September 1984.
32. Ridlen RL, Castro C, Rollheiser JJ, Kuhn GO, **Jameson CW**. Development of Analytical Methods for a Referee Dosage Analysis Program. Proceedings of the XI Federation of Analytical Chemistry and Spectroscopy Societies Meeting, Philadelphia, PA, September 1984.
33. **Jameson CW**, Goehl TJ, Davies CL. Chemical Quality Assurance Techniques for Toxicity Testing of Environmental Chemicals. Proceedings of the 1984 International Chemical Congress of Pacific Basin Societies, American Chemical Society, Honolulu, HI, December 1984.
34. Melnick RL, **Jameson CW**, Goehl TJ, Collins BJ, Maronpot RR, Greenwell A, Harrington F, Wilson R, Tomaszewski K, Agarwal D. Microencapsulation of Trichloroethylene: Stability, Bioavailability and Toxicity. North Carolina Chapter of the Society of Toxicology Annual Meeting, Research Triangle Park, NC, February 1985.
35. Dieter MP, **Jameson CW**, French JE, Brown RD, Lilya H. Target Organ Accumulation and Toxicity of Titanocene Dichloride (TDC) in F344 Rats. *The Toxicologist*, 5, 89, 1985.
36. Melnick RL, Goehl TJ, Collins B, **Jameson CW**, Maronpot R, Greenwell A, Harrington F, Wilson R, Tomaszewski K, Agarwal D. Toxicity of Microencapsulated Trichloroethylene (TCE) in Rats. *The Toxicologist*, 5, T228, 1985.
37. Kuhn GO, Dux TP, Ridlen RL, Stelting KM, **Jameson CW**. Analytical Procedures for Microencapsulated Trichloroethylene (TCE) In Rodent Feed. Proceedings of the 191st National Meeting of the American Chemical Society, Chicago, IL, September 1985.
38. Thigpen JE, Li LA, Richter CB, Lebetkin EH, Haseman JK, **Jameson CW**. The Comparative Estrogenic Activity of Semipurified, Certified, Standard and Open Formula Rodent Diets. American Association for Laboratory Animal Science, Baltimore, MD, November 1985.
39. **Jameson CW**, Schwetz BA, Goehl TJ, Moorman CR, Collins BJ. Lack of Evidence for Involvement of Cyanide in Methyl Isocyanate (MIC) Toxicity. *The Toxicologist*, 6, 76, 1986.
40. Dieter MP, **Jameson CW**, Tucker AN, Luster MI, French JE, Boorman GA. Immunotoxic Effects of Nickel Sulfate in Mice. *The Toxicologist*, 6, 263, 1986.
41. Ridlen RL, Kuhn GO, Arneson DW, **Jameson CW**. Effect of Microencapsulation Processes on Chemicals for Toxicology Studies. *The Toxicologist* 6, 304, 1986.
42. **Jameson CW**, Moseman RF, Collins BJ, Hooper ND. Spy Dust: Detection and Cleanup. Proceedings of the 193rd National Meeting of the American Chemical Society, Anaheim, CA, September 1986.
43. Moseman RF, **Jameson CW**, Hooper ND, Collins BJ. Analysis and Stability of 5-(4-Nitrophenyl)-pentadienal (NPPD) in Various Sample Materials. Proceedings of the 193rd National Meeting of the American Chemical Society, Anaheim, CA, September 1986.

C W Jameson - Curriculum Vitae and Bibliography

44. Dieter MP, French JE, **Jameson CW**, Stefanski SA, Chhabra RS. Evaluation of Mononuclear Cell Leukemia Using a Cellular Transplant Model in Rats: Studies of Pyridine, Trichlorophenol, Ethoxyethanol and Hexylresorcinol. *J. Cell. Biochem.*, 12C, 86, 1988.
45. **Jameson CW**, Goehl TJ, Gorski T, Collins BJ, Melnick RN, Kuhn GO, Harrington F. Microencapsulation of 1,1,1-Trichloroethane (TCE) and 2-Ethylhexanol (2 EH) for Toxicity Studies; Stability, Palatability and Evaluation of Chemical Availability. *The Toxicologist*, 8, 264, 1988.
46. Dieter MP, **Jameson CW**, Lodge JW, Hejtmancik M, Grumbein SL, Peters. Comparative Toxicity and Tissue Distribution of Antimony Potassium Tartrate in Rats and Mice Dosed by Drinking Water (DW) or Intraperitoneal Injection (IP). *The Toxicologist*, 8, 20, 1988.
47. Abdo KM, Thompson M, Bucher J, **Jameson CW**, Seely JC, Seilkop S, Prillaman J. Influence of Diet and Route of Administration on Vitamin E Toxicity in Male 344 Rats. *FASEB Journal*, V2, N5, 1097, 1988.
48. Yang RSH, Goehl TJ, **Jameson CW**, Germolec D, Luster MI, Chapin R, Morressey RE, Schwetz BA, Harris R, Chatham A, Arneson DW, Moseman R, Collinsworth N, Bigelow D. Chemical Stability and Biological Suitability of a 25-Chemical Mixture of Groundwater Contaminants for Animal Toxicology Studies. *The Toxicologist*, 9, 216, 1989.
49. Dieter MP, **Jameson CW**, French JE, Maronpot RR, Chhabra RS, Chan PC. Investigation of Structure Activity Relationships Between NTP Test Chemicals and Mononuclear Cell Leukemia Using a Cellular Transplant Model. *The Toxicologist*, 9, 124, 1989.
50. Michael LC, Martin-Goldberg M, Handy RW, **Jameson CW**. Determination of Lead and Cadmium in Rat Liver by Square Wave Stripping Voltammetry. *Pittsburgh Conference*, 1989.
51. Dieter MP, **Jameson CW**, Maronpot RR, Braun AG. The Chemotherapeutic Potential of Glycol Ethers. Structure Activity Studies with a Cellular Transplant Model of Leukemia in Rats. *Proc. Amer. Assoc. Cancer Res.*, 30, 615, 1989.
52. Bursey J, Moseman R, Parker C, Tomer K, **Jameson CW**. GC/MS and LC/MS in the Analysis of NPPD (Spy Dust). *Proceedings of the Annual Meeting of The American Society of Mass Spectrometry*, 1989.
53. Dieter MP, **Jameson CW**, Maronpot RR, French JE. Utilization of a Fischer Rat Leukemia Transplant Model as a Short-Term Assay to Screen Chemicals for Leukemogenesis or for Potential Anti-Leukemic Activity. *Proceedings, XIVth International Symposium for Comparative Research on Leukemia and Related Disease*, 1989.
54. Dieter MP, **Jameson CW**, Elwell MR. Structural Alerts for Leukemia: The Alkyl Phosphonic Ester (APE) Structure. *Proc. Amer. Assoc. Cancer Res.*, 1990.
55. **Jameson CW**, Eustis SL, Hong LH. Toxicology and Carcinogenicity Studies of d-Limonene in Male and Female F344 Rats and B6C3F1 Mice. *The Toxicologist*, 10, 142, 1990.

C W Jameson - Curriculum Vitae and Bibliography

56. Yuan J, **Jameson CW**, Goehl TJ, Collins BJ, Corniffee G. Systemic Availability of o-Nitroanisole in Male F344 Rats from Aged and Fresh Feed Formulations. *The Toxicologist*, 10, 337, 1990.
57. Dieter MP, Matthews HB, **Jameson CW**, Jeffcoat AR. Comparative Metabolism of 2,4- and 2,6-Isomers of Toluene Diisocyanate in F344 Rats. *The Toxicologist*, 10, 334, 1990.
58. Graves S, LeMunyon JM, Hardbeck EC, Arnold WT, Karrenbrock AH, Arneson DW, **Jameson CW**. Dose Formulation Studies of Microencapsulated 1,1,2,2,-Tetrachloroethane. *The Toxicologist*, 10, 255, 1990.
59. Abu-Shakra A, Johnson L, Earley K, Gupta R, **Jameson CW**, Kari F, Langenbach R. Separation of the Mutagenic Components from the Carcinogen HC Blue 1. *Proc. Environ. Mutagen. Soc. Annual Meeting*, 1990.
60. **Jameson CW**, Dieter MP, Maronpot RR, French JE. "Application of a Fischer Rat Leukemia Transplant Model as a Screen for the Leukemogenic Potential of Chemicals". *Proc. of The International Symposium on Toxicology, Beijing, P. R. China, October 16-19, 1990.*
61. Elwell MR, **Jameson CW**, Thompson MB, Yuan J, Carleton T. Toxicity of 4-Chloro- α,α,α -trifluorotoluene. *Proc. American College of Veterinary Pathologists Annual Meeting*, 1990.
62. Yuan J, **Jameson CW**, Goehl TJ, Elwell MR, Thompson MB, Corniffe G, Carleton T. Application of Molecular Encapsulation for Toxicology Studies I: Molecular Encapsulation of 4-Chloro- α,α,α -trifluoro-toluene with β $\square$ Cyclodextrin. *The Toxicologist*, 11, 1991.
63. **Jameson CW**, Yuan J, Goehl TJ, Elwell MR, Thompson MB, Corniffe G, Carleton T. Application of Molecular Encapsulation for Toxicology Studies II: 14 Day Toxicity Study of 4-Chloro- α,α,α -trifluoro-toluene in Male and Female F344 rats and B6C3F1 Mice Using β Cyclodextrin and Corn Oil Vehicles. *The Toxicologist*, 11, 1991.
64. Moseman RF, Brink RE, **Jameson CW**, Treinen KA, Chapin RE. Methods Development and Validation for the Determination of ppm Levels of Boron in Rat Tissues. *The Toxicologist*, 11, 1991.
65. Dixon DB, Moseman RF, **Jameson CW**, DeMartin FL, Heil TP, Foster SC. Method Development and Validation for the Determination of ppm Levels of Hildebrandt Acid in Blood. *The Toxicologist*, 11, 1991.
66. **Jameson CW**, Elwell M, Eustis SL, Hong LH. Toxicology and Carcinogenicity Studies of d- Limonene in Male and Female F344 Rats and B6C3F1 Mice. *Proceedings of the 204th National Meeting of the American Chemical Society, Washington, DC, August 23-28, 1992.*
67. **Jameson CW**, Lucier GW, Barrett J C. The Biennial Report On Carcinogens (BRC): Review and Revision of the Criteria for Listing Substances in the BRC. *The Toxicologist*, 30, No.1, 1996.

C W Jameson - Curriculum Vitae and Bibliography

68. Mahaffey K, Gulson B, **Jameson CW**, Vidal M, Law A, Mizon K, Smith A, Korsch M. Dietary Lead Intakes For Mother-Child Pairs And Relevance To Pharmacokinetic Models. *The Toxicologist*, 42, No.1-S, 1998.
69. Gulson B, Mahaffey K, **Jameson CW**, Mizon K, Korsch M, Cameron M, Eisman, J. Lead Mobilisation From The Maternal Skeleton Is Greater During The Post-Pregnancy Period Than During Pregnancy. *The Toxicologist*, 42, No.1-S, 1998.
70. Gulson BL, Mizon KJ, Patison N, Law AJ, Korsch MJ, **Jameson CW**, Mahaffey KR, Donnelly J. Contribution Of Lead From Formula And Food To Lead In Blood And Urine Of Newly Born Infants And Relationship To Skeletal Mobilization. *The Toxicologist*, 48, No.1-S, 1999.
71. **Jameson CW**. Report on Carcinogens: History and Process. Abstracts of Papers of the American Chemical Society, 228: U303-U303, 2004.

EXHIBIT B

Additional References

1. Robin Mesnage , George Renney , Gilles-Eric Séralini , Malcolm Ward & Michael N. Antoniou. (2017) Multiomics reveal non-alcoholic fatty liver disease in rats following chronic exposure to an ultra-low dose of Roundup herbicide. Scientific Reports (www.nature.com/scientificreports) | 7:39328 | DOI: 10.1038/srep39328
2. Lisbeth E. Knudsen, Pernille Winton Hansen, Seher Mizrak, Heidi K. Hansen, Thit A. Morck, Flemming Nielsen, Volkert Siersma and Line Mathiesen. (2017) Biomonitoring of Danish school children and mothers including biomarkers of PBDE and glyphosate. Rev Environ Health 2017; aop DOI 10.1515/reveh-2016-0067
3. Niall Shanks, Ray Greek and Jean Greek.(2019) Are animal models predictive for humans? Philosophy, Ethics, and Humanities in Medicine, 4:2 doi:10.1186/1747-5341-4-2
4. Timothy L Lash (2007). Bias analysis applied to Agricultural Health Study publications to estimate non-random sources of uncertainty. Journal of Occupational Medicine and Toxicology, 2:15
5. Gilles-Eric Seralini, Robin Mesnage, Nicolas Defarge, and Joel Spiroux de Vendomois (2014) Conflicts of interests, confidentiality and censorship in health risk assessment: the example of an herbicide and a GMO. Environmental Sciences Europe 26:13
6. EPA, R.E.D. Facts sheet: Glyphosate (1993) EPA-738-F-93-011
7. EPA, Reregistration Eligibility Decision (RED); Glyphosate, (1993) EPA 738-R-93-014
8. EPA, Glyphosate; Pesticide Tolerances, Federal Register: April 11, 1997 (Volume 62, Number 70)
9. EPA, Glyphosate Final Work Plan (WFP) Registration Review Case No. 0178, December 2009
10. EPA, Glyphosate Summary Document Registration Review: Initial Docket, June 2009
11. David Brusick, Marilyn Aardema, Larry Kier, David Kirkland & Gary Williams (2016) Genotoxicity Expert Panel review: weight of evidence evaluation of the genotoxicity of glyphosate, glyphosate-based formulations, and aminomethylphosphonic acid. CRITICAL REVIEWS IN TOXICOLOGY, 2016 VOL. 46, NO. S1, 56-74

12. Keith Fluegge (2016) Overlooking relevant confounders in the assessment of pesticides and human health: a reply to Mostafalou and Abdollahi. Arch Toxicol. Published on line December 28, 2016.
13. John Acquavella, David Garabrant, Gary Marsh, Tom Sorahan & Douglas L. Weed (2016) Glyphosate epidemiology expert panel review: a weight of evidence systematic review of the relationship between glyphosate exposure and non-Hodgkin's lymphoma or multiple myeloma. CRITICAL REVIEWS IN TOXICOLOGY, 2016 VOL. 46, NO. S1, 28-43
14. Donna R. Farmer, Timothy L. Lash, John F. Acquavella (2005) Glyphosate Results Revisited. Environmental Health Perspectives/Correspondence • VOLUME 113 | NUMBER 6 | June 2005
15. Anneclaire J. De Roos and Megan A. Svec, Aaron Blair, Jennifer A. Rusiecki, Mustafa Dosemeci, and Michael C. Alavanja, Jane A. Hoppin and Dale P. Sandler (2005) Glyphosate Results Revisited: De Roos et al. Respond Environmental Health Perspectives/Correspondence • VOLUME 113 | NUMBER 6 | June 2005
16. Jose V. Tarazona, Daniele Court-Marques, Manuela Tiramani, Hermine Reich, Rudolf Pfeil, Frederique Istace, Federica Crivellente Glyphosate toxicity and carcinogenicity: a review of the scientific basis of the European Union assessment and its differences with IARC. Arch Toxicol. Published on line December 28, 2016.
17. Channa Jayasumana, Sarath Gunatilake, and Priyantha Senanayake, Glyphosate, Hard Water and Nephrotoxic Metals: Are They the Culprits Behind the Epidemic of Chronic Kidney Disease of Unknown Etiology in Sri Lanka? Int. J. Environ. Res. Public Health 2014, 11, 2125-2147.
18. Carcinogenicity of tetrachlorvinphos, parathion, malathion, diazinon, and glyphosate. www.thelancet.com/oncology Vol 16 May 2015
19. Tom Sorahan. Multiple Myeloma and Glyphosate Use: A Re-Analysis of US Agricultural Health Study (AHS) Data. Int. J. Environ. Res. Public Health 2015, 12, 1548-1559
20. John Peterson Myers, Michael N. Antoniou, Bruce Blumberg, Lynn Carroll, Theo Colborn, Lorne G. Everett, Michael Hansen, Philip J. Landrigan, Bruce P. Lanphear, Robin Mesnage, Laura N. Vandenberg, Frederick S. vom Saal, Wade V. Welshons and Charles M. Benbrook Concerns over use of glyphosate-based herbicides and risks associated with exposures: a consensus statement. Environmental Health (2016) 15:19
21. NTP: Glyphosate Research Scoping (2016). Downloaded from: https://ntp.niehs.nih.gov/ntp/about_ntp/bsc/2016/june/presentations/glyphosate_508.pdf

22. NTP Glyphosate and Glyphosate Formulations Research (2017) Downloaded from: <https://ntp.niehs.nih.gov/results/areas/glyphosate/index.html>
23. Toxicology and Carcinogenesis Studies of Vinylidene Chloride (CAS No. 75-35-4) in F344/N Rats and B6C3F1/N Mice (Inhalation Studies). Technical Report Series No. 582. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Research Triangle Park, NC., and Bethesda, MD.
24. John Acquavella, Donna Farmer, and Mark R. Cullen (1999) A Case–Control Study of Non-Hodgkin Lymphoma and Exposure to Pesticides. *CANCER/Correspondence* August, 15 1999 / Volume 86 / Number 4
25. Lennart Hardell, Mikael Eriksson (1999) A Case–Control Study of Non-Hodgkin Lymphoma and Exposure to Pesticides - Author Reply. *CANCER/Correspondence* August, 15 1999 / Volume 86 / Number 4
26. Shahla Hosseini Bai1 & Steven M. Ogbourne1(2016) Glyphosate: environmental contamination, toxicity and potential risks to human health via food contamination. *Environ Sci Pollut Res* (2016) 23:18988–19001
27. Goen, T., et al., Efficiency control of dietary pesticide intake reduction by human biomonitoring. *Int. J. Hyg. Environ. Health* (2016),
28. Pamela J. Mink, Jack S. Mandel, Bonnielin K. Scurman, Jessica I. Lundin Epidemiologic studies of glyphosate and cancer: A review *Regulatory Toxicology and Pharmacology* 63 (2012) 440–452
29. Lennart Hardell, Mikael Eriksson & Marie Nordström (2002) Exposure to Pesticides as Risk Factor for Non-Hodgkin's Lymphoma and Hairy Cell Leukemia: Pooled Analysis of Two Swedish Case-control Studies, *Leukemia & Lymphoma*, 43:5, 1043-1049
30. Celia P.M. Bento, Dirk Goossens, Mahrooz Rezaei, Michel Riksen, Hans G.J. Mol, Coen J. Ritsema, Violette Geissen. Glyphosate and AMPA distribution in wind-eroded sediment derived from loss soil *Environmental Pollution* 220 (2017) 1079-1089.
31. Kathryn Z. Guyton, Dana Loomis, Kurt Straif Reply to “the critical role of pre-publication peer review—a case study of glyphosate” by FN Dost *LETTER TO THE EDITOR/ Environ Sci Pollut Res* published on line 10 October 2016
32. Jasmine Jomichen, Sonia El-Zaemey, Jane S Heyworth, Renee N Carey, Ellie Darcey, Alison Reid, Deborah C Glass, Tim Driscoll, Susan Peters, Michael Abramson3 Lin Fritschi Australian work exposures studies: occupational exposure to pesticides. *Occup Environ Med* 2017;74:46–51

33. Christina Avgerinou, Ioanna Giannezi, Stela Theodoropoulou, Vasileios Lazaris, Georgia Kolliopoulou, Panagiotis Zikos, Yannis Alamanos, Michalis Leotsinidis & Argiris Symeonidis (2017): Occupational, dietary, and other risk factors for myelodysplastic syndromes in Western Greece, Hematology, DOI: 10.1080/10245332.2016.1277006
34. Cristina Fortes, Simona Mastroeni, Marjorie M. Segatto, Clarissa Hohmann, Lucia Miligi, Lucio Bakos, and Renan Bonamigo. Occupational Exposure to Pesticides With Occupational Sun Exposure Increases the Risk for Cutaneous Melanoma. JOEM Volume 58, Number 4, April 2016
35. Robert E. Tarone. On the International Agency for Research on Cancer classification of glyphosate as a probable human carcinogen. European Journal of Cancer Prevention 2016
36. Gemma Arjo, Manuel Portero, Carme Pinol, Juan Vinas, Xavier Matias-Guiu, Teresa Capell, Andrew Bartholomaeus, Wayne Parrott, Paul Christou. Plurality of opinion, scientific discourse and pseudoscience: an in depth analysis of the Seralini et al. study claiming that Roundup™ Ready corn or the herbicide Roundup™ cause cancer in rats. Transgenic Res (2013) 22:255–267
37. R. Mesnage, N. Defarge, J. Spiroux de Vendomois, G.E. Seralini. Potential toxic effects of glyphosate and its commercial formulations below regulatory limits Food and Chemical Toxicology 84 (2015) 133-153
38. Seralini GE, Clair E, Mesnage R, Gress S, Defarge N, Manuela Malatesta M et al. (2014). Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. Environmental Sciences Europe, 26(1):1–14
39. Jacob GS, Garbow JR, Hallas LE, Kimack NM, Kishore GM, Schaefer J (1988). Metabolism of glyphosate in Pseudomonas sp. strain LBr. Appl Environ Microbiol, 54(12):2953–8
40. Tomlin CDS, editor (2000). The pesticide manual: a world compendium. 12th ed. Croydon: British Crop Protection Council. Available from: <http://trove.nla.gov.au/work/6273016>.
41. Gary M. Williams, Robert Kroes, and Ian C. Munro. Safety Evaluation and Risk Assessment of the Herbicide Roundup¹ and Its Active Ingredient, Glyphosate, for Humans. Regulatory Toxicology and Pharmacology 31, 117–165 (2000)
42. Scott Weichenthal, Connie Moase, and Peter Chan. A Review of Pesticide Exposure and Cancer Incidence in the Agricultural Health Study Cohort Environmental Health Perspectives • volume 118 | number 8 | August 2010

43. Scot L. Eustis, James R. Hailey, Gary A. Boorman and Joseph K. Haseman. The Utility of Multiple-Section Sampling in the Histopathological Evaluation of the Kidney for Carcinogenicity Studies. *Toxicologic Pathology* Vol. 22, No. 5, 1994
44. Townsend, M., Peck, C., Meng, W., Heaton, M., Robison, R., O'Neill, K., Evaluation of various glyphosate concentrations on DNA damage in human Raji cells and its impact on cytotoxicity, *Regulatory Toxicology and Pharmacology* (2017), doi: 10.1016/j.yrtph.2017.02.002.
45. Isis Coalova, María del Carmen Ríos de Molina, Gabriela Chaufan. Influence of the spray adjuvant on the toxicity effects of a glyphosate formulation. *Toxicology in Vitro* 28 (2014) 1306–1311
46. Lei Luo, Fei Wang, Yiyuan Zhang, Ming Zeng, Caigao Zhong & Fang Xiao (2017): In vitro cytotoxicity assessment of roundup (glyphosate) in L-02 hepatocytes, *Journal of Environmental Science and Health, Part B*, DOI: 10.1080/03601234.2017.1293449
47. Gary L. Diamond Effects of Surfactants on the Toxicity of Glyphosate, with Specific Reference to RODEO. SERA TR 97-206-1b February 6, 1997
48. Robert Owagboriaye, Gabriel Dedeke, Kehinde Ademolu, Olarenwaju Olujimi, Joseph Ashidi, Aladesida Adeyinka (2017): Reproductive toxicity of Roundup herbicide exposure in male albino rat. *Experimental and Toxicologic Pathology*, <http://dx.doi.org/10.1016/j.etp.2017.04.007>
49. Karunanayake CP, Spinelli JJ, McLaughlin JR, Dosman JA, Pahwa P, McDuffie HH (2012). Hodgkin lymphoma and pesticides exposure in men: a Canadian case-control study. *J Agromed*, 17(1):30–9.
50. Brown LM, Blair A, Gibson R, Everett GD, Cantor KP, Schuman LM et al. (1990). Pesticide exposures and other agricultural risk factors for leukemia among men in Iowa and Minnesota. *Cancer Res*, 50(20):6585–91.
51. Lee WJ, Lijinsky W, Heineman EF, Markin RS, Weisenburger DD, Ward MH (2004). Agricultural pesticide use and adenocarcinomas of the stomach and esophagus. *Occup Environ Med*, 61(9):743–9.
52. Carreon T, Butler MA, Ruder AM, Waters MA, Davis- King KE, Calvert GM et al. ; Brain Cancer Collaborative Study Group (2005). Gliomas and farm pesticide exposure in women: The Upper Midwest Health Study. *Environ Health Perspect*, 113(5):546–51.
53. Lee WJ, Colt JS, Heineman EF, McComb R, Weisenburger DD, Lijinsky W et al. (2005). Agricultural pesticide use and risk of glioma in Nebraska, United States. *Occup Environ Med*, 62(11):786–92.

54. Pahwa P, Karunanayake CP, Dosman JA, Spinelli JJ, McLaughlin JR; Cross-Canada Group (2011). Soft- tissue sarcoma and pesticides exposure in men: results of a Canadian case-control study. *J Occup Environ Med*, 53(11):1279–86.
55. Band PR, Abanto Z, Bert J, Lang B, Fang R, Gallagher RP et al. (2011). Prostate cancer risk and exposure to pesticides in British Columbia farmers. *Prostate*, 71(2):168–83
56. Kachuri L, Demers PA, Blair A, Spinelli JJ, Pahwa M, McLaughlin JR et al. (2013). Multiple pesticide exposures and the risk of multiple myeloma in Canadian men. *Int J Cancer*, 133(8):1846–58
57. Brown LM, Burmeister LF, Everett GD, Blair A (1993). Pesticide exposures and multiple myeloma in Iowa men. *Cancer Causes Control*, 4(2):153–6
58. Lee WJ, Lijinsky W, Heineman EF, Markin RS, Weisenburger DD, Ward MH (2004). Agricultural pesticide use and adenocarcinomas of the stomach and esophagus. *Occup Environ Med*, 61(9):743–9.
59. Carreon T, Butler MA, Ruder AM, Waters MA, Davis- King KE, Calvert GM et al. ; Brain Cancer Collaborative Study Group(2005). Gliomas and farm pesticide exposure in women: The Upper Midwest Health Study. *Environ Health Perspect*, 113(5):546–51.
60. Julie Marc, Odile Mulner-Lorillon, Sandrine Boulben, Dorothee Hureau, Gael Durand, Robert Belle. Pesticide Roundup Provokes Cell Division Dysfunction at the Level of CDK1/Cyclin B Activation. *Chem. Res. Toxicol.* 2002, 15, 326-331.
61. Julie Marc, Robert Belle, Julia Morales, Patrick Cormier, Odile Mulner-Lorillon. Formulated Glyphosate Activates the DNA-Response Checkpoint of the Cell Cycle Leading to the Prevention of G2/M Transition. *Toxicological Sciences*, 82, 436-442 (2004).
62. Julie Marc, Magali Le Breton, Patrick Cormier, Julia Morales, Robert Belle, Odile Mulner-Lorillon. A glyphosate-based pesticide impinges on transcription. *Toxicology and Applied Pharmacology* 203 (2005) 1-8.
63. Breanna Ford, Leslie Bateman, Leilani Gutierrez-Palominos, Robin Park, Daniel Nomura. Mapping Proteome-wide Targets of Glyphosate in Mice. *Cell Chemical Biology* (2016).
64. Maria Lioi, Maria Scarfi, Antonietta Santoro, Roccina Barbieri, Olga Zeni, Francesca Salvemini, Dino Berardino, Matilde Ursini. Cytogenetic Damage and Induction of Pro-Oxidant State in Human Lymphocytes Exposed In Vitro to Glyphosate, Vinclozolin, Atrazine, and DPX-E9636. *Environmental and Molecular Mutagenesis* 32:39-46 (1998).

65. Claudia Bolognesi, Amadeu Creus, Patricia Ostrosky-Wegman, Ricard Marcos. Micronuclei and pesticide exposure. *Mutagenesis*, vol. 26, no. 1, pp. 19-26, 2011.
66. Claudia Bolognesi, Nina Holland. The use of the lymphocyte cytokinesis-block micronucleus assay for monitoring pesticide-exposed populations. *Mutation Research* 770 (2016) 183-203.
67. Larry D. Kier, David J. Kirkland. Review of genotoxicity studies of glyphosate and glyphosate-based formulations. *Crit. Rev. Toxicol.*, 2013; 43(4): 283-315.
68. James Bus. IARC use of oxidative stress as key mode of action characteristic for facilitating cancer classification: Glyphosate case example illustrating a lack of robustness in interpretative implementation. *Regulatory Toxicology and Pharmacology* 86 (2017) 157-166.
69. Marco Peluso, Armelle Munnia, Claudia Bolognesi, Silvio Parodi. ³²P-Postlabeling Detection of DNA Adducts in Mice Treated With the Herbicide Roundup. *Environmental and Molecular Mutagenesis* 31:55-59 (1998).
70. William Heydens, Charles Healy, Kathy Hotz, Larry Kier, Mark Martens, Alan Wilson, Donna Farmer. Genotoxic Potential of Glyphosate Formulations: Mode-of-Action Investigations. *J. Agric. Food Chem.* 2008, 56, 1517-1523.
71. MONGLY01767038
72. MONGLY00586054
73. MONGLY00593610
74. MONGLY01830003 – MONGLY01830100
75. Deposition Transcripts and Exhibits of Donna Farmer, Jan 11-12, 2017.
76. Deposition Transcripts and Exhibits of Daniel Goldstein, Jan 18, 2017.
77. Deposition Transcripts and Exhibits of William Heydens, Jan. 23-24, 2017.
78. Deposition Transcripts and Exhibits of David Saltmiras, Jan 31- Feb 2, 2017.
79. Deposition Transcript and Exhibits of Aaron Blair, March 20, 2017.
80. Deposition Transcript and Exhibits of Mark Martens, April 7, 2017.

